

Increase in resting heart rate over 10-year period linked with increased risk of heart disease death

December 20 2011

In a study that enrolled nearly 30,000 apparently healthy men and women, those who had an increase in their resting heart rate over a 10-year period had an increased risk of death from all causes and from ischemic heart disease, according to a study in the December 21 issue of *JAMA*.

Some evidence indicates that a high resting heart rate (RHR) is associated with increased [cardiovascular disease](#) and death in the general population, independent of conventional [risk factors](#). However, whether changes in RHR over time influence the risk of death from ischemic [heart disease](#) (IHD) is not known, according to background information in the article.

Javaid Nauman, Ph.D., of the Norwegian University of Science and Technology, Trondheim, Norway, and colleagues conducted a study to examine the association of changes in RHR with the risk of death from IHD in a population-based group consisting of 13,499 men and 15,826 women without known cardiovascular disease. Resting [heart rate](#) was measured on two occasions around 10 years apart, with the second RHR measurement taking place between August 1995 and June 1997. There was follow-up until December 2008. A total of 60 participants were lost to follow-up due to emigration from Norway.

During an average of 12 years of follow-up, a total of 3,038 people died.

Among all deaths, 975 were caused by cardiovascular disease and 388 were due to IHD. The researchers found that compared with participants with a RHR of less than 70 beats/min at both [measurements](#), participants with a RHR of less than 70 beats/min at the first measurement but greater than 85 beats/min at the second measurement had a 90 percent increased risk of death from IHD. Participants with RHRs between 70 and 85 beats/min at the first measurement and greater than 85 beats/min at the second measurement had an 80 percent increased risk.

The researchers also found that the association of changes in RHR with all causes of death were similar to those observed for IHD mortality, but the estimates of effect were generally weaker.

Analysis also suggested that a decrease in RHR showed no general benefit in relation to IHD mortality.

"As expected from the good general health of the study participants, the observed moderate-to-strong increases in relative risk corresponded to small risk increases in absolute terms. However, it is not clear to what extent we can extrapolate our findings to less healthy individuals in whom the underlying risk is likely to be higher," the authors note.

"Our findings provide further support for the hypothesis that RHR may be an important prognostic marker for IHD and total [mortality](#). Information on RHR and its time-related changes are easy to obtain and follow-up and may be useful in identifying asymptomatic people who could benefit from measures of primary prevention, but further study in this area is warranted."

More information: *JAMA*. 2011;306[23]:2579-2587.

Provided by JAMA and Archives Journals

Citation: Increase in resting heart rate over 10-year period linked with increased risk of heart disease death (2011, December 20) retrieved 20 September 2024 from <https://medicalxpress.com/news/2011-12-resting-heart-year-period-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.