

Study: One-fifth of spine surgery patients develop PTSD symptoms

September 28 2012

Nearly 20 percent of people who underwent low back fusion surgery developed post-traumatic stress disorder symptoms associated with that surgery, according to a recent Oregon Health & Science University study published in the journal *Spine*.

Past studies have noted PTSD [symptoms](#) in some trauma, cancer and organ transplant patients. But this is the first study, its authors believe, to monitor for PTSD symptoms in patients undergoing an elective medical procedure.

"It is maybe not surprising that significant surgical interventions have psychological as well as physical impacts," said Robert Hart, M.D., an orthopedic surgeon, professor of orthopedics and rehabilitation at OHSU and senior author of the study. "I think it means that we in the medical community need to monitor for these effects in our patients and to manage them when they occur."

The study found that the strongest factor in determining whether a spine surgery patient suffered PTSD symptoms after the surgery was whether the patient had a psychiatric disturbance—depression or anxiety disorders, for example—before the surgery. But some patients who had no such pre-surgery diagnoses also suffered PTSD symptoms after the spine surgery, the study found.

The OHSU study involved 73 of Hart's surgical patients who underwent lumbar spinal fusion surgery, a procedure in which two or more lumbar

vertebrae in the back are fused together, usually including implantation of metal screws and rods as part of the procedure.

Surgeons at the OHSU Spine Center always look for other ways to deal with back pain before surgery. But sometimes, surgery is the only option. And spinal fusion is often major surgery that may involve complications and often involves a long recuperation period. The majority of patients in the study reported a good experience and had substantial relief of their pre-operative pain.

Hart said the study results might offer guidance to doctors and surgeons in advising potential [spine surgery](#) patients about the surgery. For patients at higher risk, the results suggest doctors and surgeons might offer treatment for them prior to surgery, Hart said.

"At the end of the day, I hope this will make changes in the way we prepare people for surgery, to reduce these psychological impacts. This is analogous to how we currently optimize [patients'](#) physical condition before major surgery," Hart said.

Hart said he wants to study that issue next—and measure which pre-[surgery](#) interventions might work best to reduce or eliminate any PTSD effects.

Provided by Oregon Health & Science University

Citation: Study: One-fifth of spine surgery patients develop PTSD symptoms (2012, September 28) retrieved 15 September 2024 from <https://medicalxpress.com/news/2012-09-one-fifth-spine-surgery-patients-ptsd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.