

## How to treat and not treat a cold

January 7 2013, by Randy Dotinga

---

A sniffle here, a cough there and suddenly a cold has come. What to do? Here's how to treat - and not treat - a cold.

Q: How do you know if you have a cold?

A: The "common cold" lasts three to 10 days, though a cough can stick around longer, said Dr. Donna Meltzer, residency director at Stony Brook University's Department of [Family Medicine](#).

"It is characterized by variable degrees of sore throat, sneezing, [nasal congestion](#) and runny nose, low-grade fever, generalized malaise and sometimes headache," she said. "There are many types of viruses that cause colds, and infection is spread through the air when someone coughs or [sneezes](#), or by direct contact with an infected person [shaking hands] or a contaminated surface [like a doorknob]."

Q: Are there effective home remedies?

A: Doctors still recommend the age-old standbys: Get rest and drink a lot of fluids, which help thin the mucus.

Gargling with salt water is also a good idea, said Meltzer, who recommends mixing teaspoon of salt in 8 ounces of water. Though studies differ on whether it's effective, she said that, in her experience, it temporarily relieves the pain of a sore throat.

It might even be helpful to gargle every day when you aren't sick. "Some

newer studies suggest that individuals who gargle three times daily with water have fewer [cold symptoms](#)," she said.

Q: What are the best options at the drugstore?

A: A pharmacist can help people with colds choose the best over-the-counter medication for their particular symptoms. But, it's important to tell the pharmacist about any conditions such as diabetes, [heart problems](#), [high blood pressure](#) or glaucoma because cold medications can exacerbate them, said Mark Macchia, associate director of pharmaceutical services at Mercy Medical Center in Rockville Centre.

Also note other medications that you're taking, keeping in mind that your pharmacist may not have a complete picture of your medical history, he said.

Q: What about decongestants and antihistamines?

A: "In adults, over-the-counter decongestants taken by mouth like Sudafed and [nasal sprays](#) like Neo-Synephrine might temporarily provide relief of nasal stuffiness," Meltzer said. "Studies indicate that antihistamines such as Benadryl are ineffective alone, but might relieve some nasal symptoms when combined with a decongestant. Some feel that guaifenesin for cough might help." Prescription nasal sprays may also provide relief, she said.

Macchia noted, though, that it's important to follow the instructions about how long you should use medicated nasal sprays, which can be effective in the short term but cause problems if taken over more than a few days.

Saline nasal sprays, which moisten the nasal passages, can be used indefinitely, but medicated sprays can cause what's called a rebound

effect, actually making congestion worse if they're used too long, he said. "There are more incidents of rebound congestion with the 4-hour nasal sprays," Macchia said. "The long-acting ones have fewer incidents." His recommendation for someone who wants to use a nasal spray is a 12-hour spray.

Q: Are antibiotics an option?

A: Antibiotics are not recommended unless you have a bacterial infection - and most colds are not caused by bacteria. "Patients should avoid insisting that their family physicians or pediatricians prescribe antibiotics," Meltzer said.

"Even educated patients insist on antibiotics and are convinced they work, but patients simply get better because the cold has run its course," she said. "In fact, most patients never complete the anti biotic prescription. Aside from creating resistance and cost implications, patients risk side effects."

And, she said, keep in mind that green mucus doesn't necessarily mean that you have a bacterial infection and need antibiotics.

Q: Do you treat a kid with a cold the same as an adult?

A: Avoid giving aspirin to kids because it can cause serious side effects. "In both children and adults, acetaminophen and nonsteroidal anti-inflammatory drugs such as ibuprofen are helpful in reducing a fever and relieving headache and body aches," Meltzer said.

Also keep in mind that cough and [cold medications](#) are usually not advised in children younger than 6 years and should not be used in children younger than 2, she added.

Q: How active, or inactive, should you be with a cold?

A: "If you have a fever, you should probably stay home from school and work," Meltzer said. "I also do not allow athletes to exercise or compete in sports if they have a fever."

The picture changes, though, once all that's left of your cold are what she called "residual cold symptoms" - slight cough or runny nose without fever. For the average person whose [cold](#) has progressed to that point and who feels well, she said, going to work or school is probably OK. For athletes and active adults with only "above the neck symptoms" and no fever, Meltzer said, "it is reasonable to engage in mild or moderate exercise if he or she feels well enough."

(c)2013 Newsday

Distributed by MCT Information Services

Citation: How to treat [?](#) and not treat [?](#) a cold (2013, January 7) retrieved 6 May 2024 from <https://medicalxpress.com/news/2013-01-cold.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.