

A new treatment for kidney disease-associated heart failure?

January 9 2013

Chronic kidney disease (CKD) patients frequently suffer from mineral bone disorder, which causes vascular calcification and, eventually, chronic heart failure. Similar to patients with CKD, mice with low levels of the protein klotho (klotho hypomorphic mice) also develop vascular calcification and have shorter life spans compared to normal mice.

In this issue of the *Journal of Clinical Investigation*, Florian Lang and colleagues at the University of Tübingen in Germany, found that treatment with the mineralocorticoid receptor antagonist spironolactone reduced [vascular calcification](#) in klotho hypomorphic mice and increased their life span.

In a companion Attending Physician article, Darryl Quarles of the University of Tennessee discusses the implications of these findings for the treatment of CKD patients.

More information: Spironolactone-sensitive vascular calcification and Pit-1-dependent osteoblastic differentiation in klotho-hypomorphic mice, *Journal of Clinical Investigation*, 2013.

Reducing cardiovascular mortality in chronic kidney disease: something borrowed, something new, *Journal of Clinical Investigation*, 2013.

Provided by Journal of Clinical Investigation

Citation: A new treatment for kidney disease-associated heart failure? (2013, January 9)
retrieved 23 April 2024 from

<https://medicalxpress.com/news/2013-01-treatment-kidney-disease-associated-heart-failure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.