

Treatment of asthma: Stepping up treatment and also stepping it down

April 9 2007

Asthma symptoms vary greatly among individuals and vary at times with each individual. In this comprehensive study in the April issue of *Mayo Clinic Proceedings*, researchers address the prevalence and process of step-down therapy as symptoms subside. Of the 397 adults and children studied, 64 percent had at least one change in medication dose during the two years of the study. Most changes were step-up in doses during an asthma flare. Step-down changes were far less common.

Lead author Barbara Yawn., M.D., from the Department of Research at Olmsted Medical Center in Rochester, Minn., comments that “patients indicate it is important to them to take the lowest possible dose of medication to maintain asthma control, suggesting we need to step down therapy when feasible.”

This is one of the first studies to address the frequency, type and context of step-up and step-down changes in asthma therapy for children and young adults. The study asserts that asthma care continues to be episodic, with most physician visits occurring at a flare-up. Additional research will help determine the best course of long-term treatment for patients with asthma.

Source: Mayo Clinic

Citation: Treatment of asthma: Stepping up treatment and also stepping it down (2007, April 9)

retrieved 2 May 2024 from <https://medicalxpress.com/news/2007-04-treatment-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.