

# Children of smokers have more than 5 times higher levels of a nicotine toxin

June 19 2007

---

Children who have at least one parent who smokes have 5.5 times higher levels of cotinine, a byproduct of nicotine, in their urine, according to a study by researchers from Warwick Medical School at the the University of Warwick, and the University of Leicester, published online ahead of print in *Archives of Disease in Childhood*.

Having a mother that smokes was found to have the biggest independent effect on cotinine in the urine – quadrupling it. Having a smoking father doubled the amount of cotinine, one of chemicals produced when the body breaks down nicotine from inhaled smoke to get rid of it.

Sleeping with parents and lower temperature rooms were also associated with increased amounts of cotinine.

Cotinine was measured in 100 urine samples taken from infants aged 12 weeks. Seventy one of the babies had at least one parent that smoked and the parents of the other 33 were non-smokers.

The authors say: ‘Smoking babies tend to come from poorer homes, which may have smaller rooms and inadequate heating.

‘Higher cotinine levels in colder times of year may be a reflection of the other key factors which influence exposure to passive smoking, such as poorer ventilation or a greater tendency for parents to smoke indoors in winter.’

Sleeping with a parent is a known risk factor for cot death and the authors suggest that one reason for this could be inhalation of, or closeness to clothing or other objects contaminated with, smoke particles during sleep.

Nearly 40% of under-fives are believed to be exposed to tobacco smoke at home, and smoke may be responsible for up to 6,000 deaths per year in young children.

The authors say: 'Babies and children are routinely exposed to cigarette smoking by their carers in their homes, without the legislative protection available to adults in public places.'

But they acknowledge that there are practical difficulties in preventing smoking in private homes because it relies on parents or carers being educated about the harmful effects of passive smoking on their children and then acting on that knowledge.

Source: University of Warwick

Citation: Children of smokers have more than 5 times higher levels of a nicotine toxin (2007, June 19) retrieved 6 May 2024 from <https://medicalxpress.com/news/2007-06-children-smokers-higher-nicotine-toxin.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--