

Prevent smoking to reduce risk of erectile dysfunction

July 27 2007

Men who smoke cigarettes run an increased risk of experiencing erectile dysfunction, and the more cigarettes smoked, the greater the risk, according to a study by Tulane University researchers published in the American Journal of Epidemiology.

A team of researchers led by Jiang He, Professor of Epidemiology at the Tulane University School of Public Health and Tropical Medicine, examined the association between cigarette smoking and erectile dysfunction in a 2000-2001 study in China involving 7,684 men. The researchers used questionnaires to assess the status of cigarette smoking and erectile dysfunction. Those surveyed were men between the ages of 35-74 who did not have vascular disease.

The team found that there was a significant statistical link between the number of cigarettes that men smoked and the likelihood they would experience erectile dysfunction. The association between smoking and erectile dysfunction was even stronger in participants with diabetes. An estimated 22.7 percent of erectile dysfunction cases among Chinese men might be attributable to cigarette smoking, says the study.

Although erectile dysfunction is not a life-threatening condition, it compromises well-being and quality of life. The Tulane study results suggest that smoking prevention should be an important approach for reducing the risk of erectile dysfunction.

Source: Tulane University

Citation: Prevent smoking to reduce risk of erectile dysfunction (2007, July 27) retrieved 9 April 2024 from <https://medicalxpress.com/news/2007-07-erectile-dysfunction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.