

# Hallucinations in schizophrenia linked to brain area that processes voices

July 31 2007

---

For the first time, researchers using magnetic resonance imaging (MRI) have found both structural and functional abnormalities in specific brain regions of schizophrenic patients who experience chronic auditory hallucinations, according to a study published in the August issue of Radiology.

“The results showed abnormalities in specific areas of the brain associated with the capacity to process human voices,” said lead author, Luis Martí-Bonmatí, M.D., Ph.D., chief of magnetic resonance in the Department of Radiology at Dr. Peset University Hospital in Valencia, Spain.

Schizophrenia is a chronic, severe and disabling brain disorder that affects approximately 1 percent of the global population, according to the National Institute of Mental Health. People with schizophrenia often experience hallucinations, delusional thoughts, paranoia, disorganized thinking and other cognitive difficulties. Although the cause of schizophrenia has not been determined, it is believed to result from a combination of environmental and genetic factors. The condition is treatable, but there is no cure.

“Developing a clear understanding of the pathological abnormalities associated with schizophrenia is one of the greatest challenges in psychiatry,” Dr. Martí-Bonmatí said. “Using MRI to mark brain regions that are affected in both structure and function will help pinpoint specific abnormalities associated with the disease and ultimately enable

more effective treatment.”

The researchers studied 31 right-handed men, 21 with schizophrenia who suffered from persistent auditory hallucinations and 10 healthy controls. Morphological MR images were acquired to show abnormalities in brain structure while functional MRI was used to gauge brain response to various emotional and neutral stimuli.

Among the schizophrenic patients, the results showed functional abnormalities and corresponding gray matter deficits in several brain regions associated with regulating emotion and processing human voices.

“We hope that by evaluating combined structural and functional abnormalities in the brains of these patients, we may uncover biological markers to find candidates for specific treatments and better monitor patient response to those treatments,” Dr. Martí-Bonmatí said.

Source: Radiological Society of North America

Citation: Hallucinations in schizophrenia linked to brain area that processes voices (2007, July 31) retrieved 4 May 2024 from <https://medicalxpress.com/news/2007-07-hallucinations-schizophrenia-linked-brain-area.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--