


# This is your brain on violent media

December 6 2007



The yellow area of the brain is the right lateral orbitofrontal cortex, or right lOFC, which has been previously associated with decreased control over a variety of behaviors, including reactive aggression. The first graph illustrates that as the number of violent movies watched increased, the right lOFC activity diminished. The second graph shows that when subjects watched the non-violent control clips, there were no systematic changes in the activation of this area. Credit: Columbia University Medical Center

Violence is a frequent occurrence in television shows and movies, but can watching it make you behave differently?

Although research has shown some correlation between exposure to media violence and real-life violent behavior, there has been little direct neuroscientific support for this theory until now.

Researchers at Columbia University Medical Center’s Functional Magnetic Resonance Imaging (fMRI) Research Center have shown that watching violent programs can cause parts of your brain that suppress

aggressive behaviors to become less active.

In a paper in the Dec. 5 on-line issue of *PLoS ONE*, Columbia scientists show that a brain network responsible for suppressing behaviors like inappropriate or unwarranted aggression (including the right lateral orbitofrontal cortex, or right lOFC, and the amygdala) became less active after study subjects watched several short clips from popular movies depicting acts of violence. These changes could render people less able to control their own aggressive behavior. Indeed the authors found that, even among their own subjects, less activation in this network was characteristic of people reporting an above average tendency to behave aggressively. This characteristic was measured through a personality test.

A secondary finding was that after repeated viewings of violence, an area of the brain associated with planning behaviors became more active. This lends further support to the idea that exposure to violence diminishes the brain's ability to inhibit behavior-related processing.

None of these changes in brain activity occurred when subjects watched non-violent but equally engaging movies depicting scenes of horror or physical activity.

“These changes in the brain’s behavioral control circuits were specific to the repeated exposure to the violent clips,” said Joy Hirsch, Ph.D., professor of Functional Neuroradiology, Psychology, and Neuroscience and Director of the Center for fMRI at CUMC, and the *PLoS ONE* paper’s senior author. “Even when the level of action in the control movies was comparable, we just did not observe the same changes in brain response that we did when the subjects viewed the violent clips.”

“Depictions of violent acts have become very common in the popular media,” said Christopher Kelly, the first author on the paper and a

current CUMC medical student. “Our findings demonstrate for the first time that watching media depictions of violence does influence processing in parts of the brain that control behaviors like aggression. This is an important finding, and further research should examine very closely how these changes affect real-life behavior.”

Source: Columbia University Medical Center

Citation: This is your brain on violent media (2007, December 6) retrieved 20 April 2024 from <https://medicalxpress.com/news/2007-12-brain-violent-media.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--