

Neurotransmitters in biopolymers stimulate nerve regeneration

December 11 2007



Fluorescence micrograph of a ganglion on a 70 percent acetylcholine polymer that shows neurites expressing an established neuronal marker called synaptophysin. The bright red spots on the neurites indicate the presence of synaptic vesicle proteins, which are required for functional restoration after nerve injury. Credit: Image courtesy of Christiane Gumera

Research reported December 11 in the journal *Advanced Materials* describes a potentially promising strategy for encouraging the regeneration of damaged central nervous system cells known as neurons.

The technique would use a biodegradable polymer containing a chemical



group that mimics the neurotransmitter acetylcholine to spur the growth of neurites, which are projections that form the connections among neurons and between neurons and other cells. The biomimetic polymers would then guide the growth of the regenerating nerve.

There is currently no treatment for recovering human nerve function after injury to the brain or spinal cord because central nervous system neurons have a very limited capability of self-repair and regeneration.

"Regeneration in the central nervous system requires neural activity, not just neuronal growth factors alone, so we thought a neurotransmitter might send the necessary signals," said Yadong Wang, assistant professor in the Coulter Department of Biomedical Engineering at Georgia Tech and Emory University, and principal investigator of the study. The research was supported by Georgia Tech, the National Science Foundation and the National Institute of Biomedical Imaging and Bioengineering (NIBIB).

Chemical neurotransmitters relay, amplify and modulate signals between a neuron and another cell. This new study shows that integrating neurotransmitters into biodegradable polymers results in a biomaterial that successfully promotes neurite growth, which is necessary for victims of central nervous system injury, stroke or certain neurodegenerative diseases to recover sensory, motor, cognitive or autonomic functions.

Wang and graduate student Christiane Gumera developed novel biodegradable polymers with a flexible backbone that allowed neurotransmitters to be easily added as a side chain. In its current form, the polymer would be implanted via surgery to repair damaged central nerves.

"One of our ultimate goals is to create a conduit for nerve regeneration that guides the neurons to regenerate, but gradually degrades as the



neurons regenerate so that it won't constrict the nerves permanently," explained Wang.

For the experiments, the researchers tested polymers with different concentrations of the acetylcholine-mimicking groups. Acetylcholine was chosen because it is known to induce neurite outgrowth and promote the formation and strengthening of synapses, or connections between neurons. They isolated ganglia nervous tissue samples, placed them on the polymers and observed new neurites extend from the ganglia.

Since these neuron extensions must traverse a growth inhibiting material in the body, Wang and Gumera tested the ability of the biomaterial to enhance the extension of sprouted neurites. More specifically, they assessed whether the ganglia sprouted at least 20 neurites and then measured neurite length and neurite length distribution with an inverted phase contrast microscope.

"We found that adding 70 percent acetylcholine to the polymer induced regenerative responses similar to laminin, a benchmark material for nerve culture," said Wang. Seventy percent acetylcholine also led to a neurite growth rate of up to 0.7 millimeters per day, or approximately half the thickness of a compact disc.

Laminin is a natural protein present in the nervous tissues, but it dissolves in water, making it difficult to incorporate into a conduit that needs to support nerves for months. A synthetic polymer with acetylcholine functional groups, on the other hand, can be designed to be insoluble in water, according to Wang.

Since functional restoration after nerve injury requires synapse formation, the researchers also searched for the presence of synaptic vesicle proteins on the newly formed neurites. With fluorescence imaging, they found that neurons cultured on these acetylcholine



polymers expressed an established neuronal marker called synaptophysin.

To provide insights to new approaches in functional nerve regeneration, the researchers are currently investigating the mechanisms by which the neurons interact with these polymers. Since neurons that remain intact after severe injury have only a limited capacity to penetrate the scar tissue, these new findings in nerve regeneration could help compensate for the lost connections.

"This polymer and approach aren't limited to nerve regeneration though, they can probably be used for other neurodegenerative disorders as well," added Wang.

Source: Georgia Institute of Technology

Citation: Neurotransmitters in biopolymers stimulate nerve regeneration (2007, December 11) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2007-12-neurotransmitters-biopolymers-nerve-regeneration.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.