

Most with high blood pressure do not follow recommended diet

February 11 2008

A relatively small proportion of individuals with hypertension (high blood pressure) eat diets that align with government guidelines for controlling the disease, according to a report in the February 11 issue of Archives of Internal Medicine, one of the JAMA/Archives journals. In fact, since the introduction of a diet shown to help reduce blood pressure, the dietary quality of those with hypertension has decreased.

Results of the Dietary Approaches to Stop Hypertension (DASH) trial, published in 1997, indicated that a diet high in fruits, vegetables and low-fat dairy products could significantly lower blood pressure, according to background information in the article. Subsequent studies found that this eating pattern complements other lifestyle changes, such as reducing sodium and losing weight, and can help lower cholesterol levels and prevent components of the metabolic syndrome. The DASH diet is recommended by the Joint National Committee on Prevention, Detection and Treatment of High Blood Pressure for all patients with hypertension.

Philip B. Mellen, M.D., M.S., of the Hattiesburg Clinic, Hattiesburg, Miss., and colleagues analyzed data from the National Health and Nutrition Examination Survey (NHANES), conducted by the Centers for Disease Control and Prevention in 1988 to 1994 and 1999 to 2004. Men and women age 20 years and older were interviewed to obtain demographic and socioeconomic information and diet history; their blood pressure was measured, and they were asked if they had ever been diagnosed with hypertension. Using the DASH guidelines, the researchers identified goals for eight target nutrients: total fat, saturated

fat, protein, fiber, cholesterol, calcium, magnesium and potassium. DASH scores ranging from zero to nine were calculated for individuals with and without hypertension from the surveys; those who scored a 4.5 or higher were considered to be following the diet.

Among individuals surveyed in 1999 to 2004, 4,386 (28 percent) had been diagnosed with hypertension. These individuals had an average DASH score of 2.92 (compared with 3.12 among those without high blood pressure), and only 19.4 percent followed the diet. Compared with those surveyed in 1988 to 1994, 7.3 percent fewer individuals with hypertension followed the DASH diet, reflecting fewer patients who consumed target levels of total fat, fiber and magnesium. DASH scores were lowest among young people, African-Americans and those with a body mass index of 30 or higher.

Source: JAMA and Archives Journals

Citation: Most with high blood pressure do not follow recommended diet (2008, February 11) retrieved 19 April 2024 from <https://medicalxpress.com/news/2008-02-high-blood-pressure-diet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.