

Low levels of PYY hormone a very early indicator of type 2 diabetes

March 10 2008

It may soon be possible to take a simple blood test and predict whether or not someone has low levels of a particular molecule, predisposing them to the development of Type 2 diabetes. If the test is positive, it may then be possible to use preventative treatment, slowing down, or even halting that development.

Such is the hope of scientists and clinicians at Sydney's Garvan Institute of Medical Research who have shown conclusively that people who produce low levels of the molecule PYY have a higher risk of developing Type 2 diabetes and obesity.

The findings were published online on 4 March in the prestigious *International Journal of Obesity*.

It is already known that the hormone PYY, which is released from the gut after a meal, creates a feeling of satiety. When PYY is in oversupply, it prevents diet-induced obesity in mice.

Professor Herbert Herzog, Director of Garvan's Neuroscience Program, and an expert on appetite, says that the new findings are important in that they show a metabolic defect before the presence of any disease or manifestation of weight gain. "We can now see that low PYY levels after eating are a very early predictor of the development of obesity and Type 2 diabetes," he said.

Professor Lesley Campbell, Director of Diabetes Services at St.



Vincent's Hospital and a senior member of Garvan's Diabetes and Obesity Clinical Studies group, has been researching genetic factors in the development of Type 2 diabetes for over 10 years. Specifically, her research looks at people before they get the disease, the contributing factors, and the effects of the diabetes.

Professor Campbell has already published findings that insulin resistant people, with a family history of Type 2 diabetes, have low levels of PYY. "In earlier studies we hinted at the fact that before any of the abnormalities of diabetes are present, people already have an abnormality of satiety, marked by the lack of the secretion of this PYY hormone," she said.

"We now have published that, even earlier in the development of diabetes, people who are not yet insulin resistant show a low secretion of PYY. They have a blunted post-meal secretion of this hormone, making them less likely to feel satiety, and more likely to gain weight."

Professor Campbell's research involved elaborate testing of two groups of people, eight in each group, over a period of two years. One group had relatives with Type 2 diabetes, the other group had no family history of the disease. The groups were matched for gender, for age and for adiposity.

"It was most important to match the groups for their fatness," said Professor Campbell. "The only difference was their relatives. You assume that they are carrying the genetic burden of diabetes, which we already know to be a reality."

"Low levels of PYY at this very early pre-diabetes stage could be used as a marker, or predictor, that Type 2 diabetes is very likely to develop."

"As a clinician, I am hopeful that it will be possible to screen extensively



in the future, and therefore stem the spread of this debilitating disease."

Source: Research Australia

Citation: Low levels of PYY hormone a very early indicator of type 2 diabetes (2008, March 10) retrieved 1 May 2024 from https://medicalxpress.com/news/2008-03-pyy-hormone-early-indicator-diabetes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.