

## **Black women have urinary incontinence less than half as often as white women**

April 22 2008

---

The good news for black women: They have less than half the chance of developing urinary incontinence as do white women, according to a new study from the University of Michigan Health System.

The bad news: When they get it, the condition tends to be worse than in white women. The amount of urine they lose during each episode of incontinence is larger, with half of black incontinent women reporting that they lose urine to the point of noticeably wetting their underwear or a pad, compared with a third of white women.

The significance: The study confirms some common beliefs, and refutes others. The medical community has long held the belief that black women don't experience a type of urinary incontinence known as "stress incontinence," in which urine is lost during activities such as exercising, coughing and laughing. In fact, the study found, black women do experience stress incontinence. The study is in the current issue of the *Journal of Urology*.

"This is a population that may have been neglected because it was believed for so long that black women did not have stress urinary incontinence," says lead author Dee E. Fenner, M.D., Furlong Professor of Women's Health, and director of gynecology, at the U-M Health System.

"In truth, black women suffer from the social embarrassment of urinary incontinence, and the medical community needs to remember this when

diagnosing and treating all women.”

The study indicates that black women experience “urge incontinence” twice as often as white women, which supports other research on the subject. This type of incontinence involves a strong and sudden need to urinate, followed by leakage.

Additionally, it has been thought in the past that the other medical conditions associated with urinary incontinence are different between black and white women. This study suggests, however, that those conditions – such as diabetes, constipation, depression, obesity and chronic lung disease – occur at similar rates between the two races.

By the numbers:

- About 27 percent of all women surveyed had the condition.
- This study found that 14.6 percent of black women and 33.1 percent of white women have urinary incontinence.
- Black women with incontinence reported having stress incontinence in about 25 percent of instances, compared with 39 percent of white women.
- Black women with incontinence reported urge incontinence in 24 percent of cases, compared with 11 percent of white women. The remaining numbers had a combination of both types.
- The women in the study ranged from 35 to 64 years old, with an average age of 42. Most of the women – nearly 70 percent – had delivered at least one baby vaginally; vaginal deliveries are often associated with urinary incontinence.
- The study involved 1,922 black women and 892 white women from three southeastern Michigan counties. Data were collected through a telephone survey.

Source: University of Michigan

Citation: Black women have urinary incontinence less than half as often as white women (2008, April 22) retrieved 19 April 2024 from <https://medicalxpress.com/news/2008-04-black-women-urinary-incontinence-white.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.