

Black patients with terminal cancer more likely to choose aggressive care at end of life

May 30 2008

Black patients with advanced cancer were more likely than whites to die in a hospital intensive care unit, reflecting a greater preference among blacks for life-extending treatment even in the face of a terminal prognosis, according to a study led by researchers at Dana-Farber Cancer Institute in Boston. The findings will be presented at the annual meeting of the American Society of Clinical Oncology in Chicago on Monday.

The report included interview data showing that blacks more often answered yes to questions such as, "Would you want the doctors here to do everything they can to keep you alive, even if you were going to die in a day or two?"

"This is the first study focused on black/white differences that prospectively asked [terminal cancer patients] what kind of care they wanted at the end of life, and then documented the kind of care they actually received and the place of their death," said Elizabeth Trice, MD, PhD, of Dana-Farber, lead author.

Although they ruled out a number of possible explanations for the black/white differences, the investigators weren't able to identify precisely why blacks tended to prefer more-aggressive care.

"There is something different about the way black patients and white patients approach the end of life," Trice said, which may be based in cultural attitudes, religious beliefs, and how thoroughly they have been informed about and comprehend their prognosis, among other things.

Data on the preferences was obtained from the Coping with Cancer study led by Holly Prigerson, PhD, director of the Center for Psychosocial Oncology and Palliative Care Research at Dana-Farber and an associate professor of psychiatry at Harvard Medical School. That study is recruiting 800 cancer patients and their informal caregivers, such as family members.

The researchers recorded the location of death for 231 white and 61 black patients who had stage IV metastatic cancer, and who had been interviewed when they entered the study. Black patients were over four times more likely to die in a hospital ICU than white patients, they found.

The researchers, using multivariable models, found that the increased likelihood of dying in the ICU for black compared to white patients was not explained by differences in education, physical or mental health, insurance, social support, doctor-patient communication, or advance care planning, when taking into account the patient's own preference for more-aggressive care. It was clear that a patient's preference for aggressive care was the strongest factor in predicting death in an ICU.

In their initial interviews, black patients reported having a higher quality of life than their white counterparts and appeared more at peace, Trice said, which could be a factor in opting for a treatment plan aimed at extending life.

Prigerson, the senior author of the study, said the crucial question to be explored is whether the treatment preferences that explain the black/white disparities in ICU death are informed preferences or not. Toward that end, Trice has developed a research tool for assessing patients' knowledge of risks and benefits of life-extending therapies. If further research determines that blacks and whites are not equally well-informed about these risks and benefits, interventions aimed at

eliminating this disparity should be considered, said the scientists.

Source: Dana-Farber Cancer Institute

Citation: Black patients with terminal cancer more likely to choose aggressive care at end of life (2008, May 30) retrieved 4 May 2024 from <https://medicalxpress.com/news/2008-05-black-patients-terminal-cancer-aggressive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.