

Premature tooth loss can affect oral health for years to come

May 22 2008

The prospect of exchanging a tooth for that coveted reward from the tooth fairy often has kids wiggling teeth with vigor, but what happens when a primary or permanent tooth is lost prematurely due to trauma?

According to a study published in the March/April 2008 issue of *General Dentistry*, the AGD's clinical, peer-reviewed journal, parents and caretakers more often than not do not know what to do with a traumatically affected tooth and do not take proper steps to respond to the injury, which can affect their child's oral health permanently.

Two kinds of traumatic tooth loss can occur during childhood: The child can lose a primary (baby) tooth prematurely, or they can lose a permanent tooth. "With primary or "baby" teeth [the mistake parents or caretakers make] is that they often don't believe that the loss of a primary tooth is especially important since a permanent tooth is supposed to come in and replace it anyway," says AGD spokesperson, Tom A. Howley, Jr., DDS, MAGD. It is important to keep baby teeth because they maintain the spacing for permanent teeth. Keeping baby teeth in place will also increase the likelihood that permanent teeth come in straight.

According to the study's lead author, Lucianne Cople Maia, DDS, MSD, PhD, another mistake parents and caretakers make is that they believe it can always be replanted. In some cases, the tooth can be splinted back in; however, treatment still should be sought for a prematurely lost primary tooth. The area where a tooth has been lost "should be checked for bone

fractures or other damage,” among other concerns, explains Dr. Howley.

When it comes to the loss of a permanent tooth, parents and caretakers often delay in seeking treatment and/or fail to store the tooth properly en route to the hospital. AGD spokesperson Mark Donald, DMD, FAGD, explains that “the chance for success is directly related to the amount of trauma and the length of time the tooth is outside of the oral cavity. The tooth should be placed in a moist solution like milk while en route to the dentist.” Dr. Maia adds that “the ideal time between tooth loss and replantation is a period no longer than 30 minutes.”

Young children tend to lose permanent teeth prematurely the most, according to the study, due to the fact their tooth roots and gums are still developing. Parents need to be aware “that as a child becomes active, the chances of trauma to teeth will increase,” says Dr. Donald. “If and when trauma to the oral cavity occurs, the parent should get their child to the dentist as quickly as possible. “As the child begins playing sports, make sure the child wears a protective mouth guard as a safeguard to traumatic blows to the teeth,” he advises.

Parents can prepare for potential dental trauma by creating an emergency care kit that includes the dentist’s phone numbers (home and office), saline solution, a handkerchief, gauze, a small container with a lid, and ibuprofen. It is not advised to include aspirin, since it can cause excessive bleeding in an emergency.

Proper response to a traumatically affected tooth:

- Remain calm and assess the injury.
- Control the bleeding.
- Determine if the injury involves more than just tooth loss and needs medical attention.
- Call and arrange for professional care by the child’s dentist.

-- If the loss tooth is a permanent tooth, first make sure it is clean by rinsing it off with milk and replant it immediately. Hold it in place en route to receiving professional care. If it is impossible to replant the tooth, ask the dentist what medium to store the tooth in. The possibilities range from water, milk or the child's saliva. Let the dentist give directions on which solution to use.

Source: Academy of General Dentistry

Citation: Premature tooth loss can affect oral health for years to come (2008, May 22) retrieved 10 April 2024 from

<https://medicalxpress.com/news/2008-05-premature-tooth-loss-affect-oral.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|