

Study finds link between amphetamine abuse and heart attacks in young adults

June 4 2008

Young adults who abuse amphetamines may be at greater risk of suffering a heart attack, UT Southwestern Medical Center researchers have found.

In the study, available online in the journal *Drug and Alcohol Dependence*, researchers examined data from more than 3 million people between 18 and 44 years old hospitalized from 2000 through 2003 in Texas and found a relationship between a diagnosis of amphetamine abuse and heart attack.

Individual case reports have suggested a link between heart attack and amphetamine abuse, but this is believed to be the first epidemiological study of a large group of people on the issue, said Dr. Arthur Westover, assistant professor of psychiatry at UT Southwestern and the study's lead author.

"Most people aren't surprised that methamphetamines and amphetamines are bad for your health," Dr. Westover said. "But we are concerned because heart attacks in the young are rare and can be very debilitating or deadly."

Amphetamines are stimulants that can be used to treat medical conditions such as attention-deficient disorder. They are illegally abused as recreational drugs or performance enhancers.

The researchers note that abuse of methamphetamine, a type of

amphetamine often sold illegally, is increasing in most major U.S. cities.

In Texas, the researchers found greater amphetamine abuse in the north and Panhandle regions.

"This paper sounds a warning to amphetamine abusers, alerts emergency department personnel to look for amphetamine abuse in young heart attack patients, and it allows us to focus preventive efforts in geographical areas where the problems are greatest," said Dr. Robert W. Haley, chief of epidemiology at UT Southwestern and senior author of the study. Dr. Haley holds the U.S. Armed Forces Veterans Distinguished Chair for Medical Research, Honoring America's Gulf War Veterans.

"We're also concerned that the number of amphetamine-related heart attacks could be increasing," Dr. Westover said. "We'd rather raise the warning flag now than later. Hopefully, we can decrease the number of people who suffer heart attacks as the result of amphetamine abuse."

Amphetamines may contribute to heart attacks by increasing heart rate and blood pressure and by causing inflammation and artery spasms that limit blood to the heart muscle. More research is needed to determine the exact mechanism of how amphetamines work on the heart, he said.

The current research could help doctors determine the cause of heart attacks in young adults, as well as treatment. Doctors recognizing an amphetamine-caused heart attack might choose not to administer a beta-blocker medication, a common treatment for heart attack, because it could interact with methamphetamine to make the heart attack worse.

The results could have broad implications in the general population, Dr. Westover said. Texas ranks 27th among all states in use of methamphetamine among 18- to 25-year-old adults, according to a 2006

government report.

"We're talking about a state that is near the middle of prevalence of methamphetamine use in the United States, so it's possible that the number of heart attacks in young adults in other states with a much higher prevalence of amphetamine abuse may be higher as well," said Dr. Westover, who is a National Institutes of Health Multidisciplinary Clinical Research Scholar at UT Southwestern.

Source: UT Southwestern Medical Center

Citation: Study finds link between amphetamine abuse and heart attacks in young adults (2008, June 4) retrieved 26 April 2024 from <https://medicalxpress.com/news/2008-06-link-amphetamine-abuse-heart-young.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.