

# **Delayed sleep phase syndrome linked to irregular menstrual cycles, premenstrual symptoms in women**

June 10 2008

---

Women with delayed sleep phase syndrome are more likely to report irregular menstrual cycles and premenstrual symptoms, according to a research abstract that will be presented on Tuesday at SLEEP 2008, the 22nd Annual Meeting of the Associated Professional Sleep Societies (APSS).

The study, authored by Kari Sveum, of Northwestern University in Chicago, Ill., focused on 13 females with delayed sleep phase syndrome and 13 healthy controls. The subjects responded to a questionnaire regarding their reproductive health, including irregularity of their cycle and premenstrual symptoms, either in the past or present.

According to the results, twice as many subjects with delayed sleep phase syndrome reported an irregular menstrual cycle compared to controls. For those not using birth control, three times as many subjects with delayed sleep phase syndrome reported irregular menstruation, compared to controls. Pre-menstrual problems, such as cramps and mood swings, were reported by 69 percent of those with delayed sleep phase syndrome, compared to 16.67 percent of controls.

“While the data is preliminary, these results suggest that women with delayed sleep phase syndrome may be at increased risk for menstrual irregularity associated with circadian misalignment,” said Sveum.

“Further investigation with a larger group of subjects using prospective

diary data would be useful to further establish the effects of circadian disruption on reproductive cycles in women with delayed sleep phase syndrome.”

Sleep plays a vital role in promoting a woman’s health and well being. Getting the required amount of sleep is likely to enhance a woman’s overall quality of life. Yet, women face many potential barriers – such as life events, depression, illness, and medication use – that can disrupt and disturb her sleep.

The hormonal and physical changes that occur during and after menopause can also affect a woman’s sleep. Sleep disturbances are more common, and sleep quality can decline. Insomnia related to menopause often occurs.

Obstructive sleep apnea (OSA) is much more common in postmenopausal women. This increase may be due in part to menopause-related weight gain. But it also appears to be hormone-related. Estrogen seems to help protect women against OSA.

It is recommended that women get between seven and eight hours of nightly sleep.

Source: American Academy of Sleep Medicine

Citation: Delayed sleep phase syndrome linked to irregular menstrual cycles, premenstrual symptoms in women (2008, June 10) retrieved 3 May 2024 from <https://medicalxpress.com/news/2008-06-phase-syndrome-linked-irregular-menstrual.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---