

# Silencing of molecular 'conversation' may help curb severe allergies

June 30 2008

---

Scientists in Sydney have identified a process, a synergistic encounter between two molecules, that may account for the extreme allergic reactions some people experience. By silencing at least one of these molecules, it may be possible to treat allergies.

The molecules, IL-4 and IL-21, are cytokines or 'chemical messengers' produced by immune cells known as T cells. T cells use cytokines to communicate with B cells, which then make antibodies. When IL-4 and IL-21 are involved in the same 'conversation' with a B cell, laboratory experiments show they stimulate the production of large amounts of the antibody class known as immunoglobulin E, or IgE.

IgE is a very tightly regulated antibody. Under normal circumstances, it is present in only very small quantities and protects us against parasites. Unfortunately, it becomes damaging when the body makes too much.

The findings, made by Danielle Avery and Dr Stuart Tangye from the Garvan Institute of Medical Research, are now published online in the prestigious international publication *Blood*.

Dr Tangye, a B cell biologist, is interested in understanding how the body regulates the production of antibodies, in this case the IgE class. "IgE is such an efficient molecule that too much of it can be dangerous," he said. "High quantities tend to over-activate other immune cells and it's the action of those other cells that can be damaging. High levels of IgE are associated with allergies including asthma, dermatitis and rhinitis."

"It's been known for many years that IL-4 can drive IgE production in humans and mice. Our finding shows that IL-21 also stimulates production of IgE by human B cells, but it does this by using an entirely different pathway."

"As it happens, the combination of IL-4 and IL-21 provokes a very strong IgE response, around ten times greater than either molecule in isolation."

"Through an ongoing collaboration with Canberra's John Curtin School of Medical Research, Canberra Hospital and Westmead Hospital, we were able to test our theory by looking at patients with mutations in the IL-21 pathway. If you can't activate that pathway, you don't get the synergistic effect between the two cytokines."

"We surmise as a result of this study that it should be possible to target the IL-21 molecule with an antibody to block its ability to activate B cells. This may prove an effective treatment in cases where allergic responses are caused by the synergistic effect of IL-4 and IL-21."

Source: Research Australia

Citation: Silencing of molecular 'conversation' may help curb severe allergies (2008, June 30) retrieved 6 May 2024 from

<https://medicalxpress.com/news/2008-06-silencing-molecular-conversation-curb-severe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--