

Strategies to control TB outdated, inadequate

August 1 2008

The standard regimens to treat tuberculosis (TB) are inadequate in countries with high rates of multi-drug resistant (MDR) TB. In countries with high rates of MDR-TB, patients are nearly twice as likely to fail their initial treatment than those in countries with low rates, according to a new analysis of World Health Organization (WHO) data. This finding suggests strongly that current TB treatment regimens need to be updated and revised to address the shifting landscape of public health in the face of MDR-TB.

"In countries with low prevalence of initial multi-drug resistance, the standardized treatment regimens for new cases appear to be adequate," wrote Dick Menzies, M.D., lead author of the study and director of the respiratory division at McGill University. "However, in countries where the prevalence of initial drug resistance exceeds three percent, we believe it is urgent to strengthen capacity to perform drug sensitivity testing, or to reevaluate these standard treatment regimens, given the unacceptably high rates of failure and relapse."

The study appeared in the first issue for August of the American Thoracic Society's *American Journal of Respiratory and Critical Care Medicine*.

When the current public health strategies to contain TB were conceived, MDR-TB was much less common. "We hypothesized that, in countries using standardized initial and retreatment regimens, the proportion of patients with poor treatment outcomes would be correlated with prevalence of initial and acquired multi-drug resistance," wrote Dr.

Menzies.

To determine if that were the case, Dr. Menzies and co-investigators reviewed the WHO's data from 2003 and 2004 for a total of 155 countries, 121 of which reported at least 250 cases annually. They assessed dropout, failure, relapse and death rates with initial treatment, as well as dropout, failure and death rates for retreatment. All rates were analyzed with respect to the prevalence of MDR-TB in each country.

They were right: rates of failed treatment were dramatically higher with increasing prevalence of MDR-TB (p

Citation: Strategies to control TB outdated, inadequate (2008, August 1) retrieved 25 April 2024 from <https://medicalxpress.com/news/2008-08-strategies-tb-outdated-inadequate.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--