

Air pollution can hinder heart's electrical functioning

September 8 2008

Microscopic particles in polluted air can adversely affect the heart's ability to conduct electrical signals in people with serious coronary artery disease, researchers reported in *Circulation*: Journal of the American Heart Association.

In a recent study of 48 Boston-area patients, all of whom had coronary artery disease, 24-hour Holter monitors were used to examine electrocardiograms for the conductivity change called an ST-segment depression, which may indicate inadequate blood flow to the heart or inflamed heart muscle.

The average 24-hour levels for all pollutants included in the analysis were below accepted or proposed National Air Quality Standard thresholds, meaning patients were breathing air considered healthy.

"We found that an elevation in fine particles, from non-traffic as well as traffic sources, and black carbon, a marker for traffic, predicted ST-segment depression," said Diane R. Gold, M.D., M.P.H., the study's senior author and an associate professor of medicine and environmental health at Harvard University in Boston, Mass. "Effects were greatest within the first month after hospitalization, and for patients who were hospitalized for a heart attack or had diabetes."

Previous studies have documented that exposure to road traffic can trigger heart attacks, and that particulate air pollution increases the risk for cardiac death or heart attack.



"When coal sales were banned in Dublin, Ireland, and black smoke concentrations declined by 70 percent within the next 72 months, cardiovascular deaths fell by 10 percent," said Gold, citing a study published in 2002.

The ST-segment changes Gold observed were not associated with symptoms in these patients, all of whom had experienced in-hospital procedures to examine or open up their coronary arteries. Nevertheless, the findings expand the evidence that air pollution can affect heart health, either through inflaming the heart muscle or through reducing blood flow to the heart. It suggests the need for greater vigilance by physicians and heart patients in the weeks after discharge from the hospital, researchers said.

The American Heart Association and the American College of Cardiology recommend that some heart patients, particularly those who have had a heart attack, delay driving for two to three weeks after leaving the hospital and avoid driving in heavy traffic because of the stress it creates.

"Our study provides additional rationale to avoid or reduce heavy traffic exposure after discharge, even for those without a heart attack, since traffic exposure involves pollution exposure as well as stress," she said.

The study's 48 participants had been hospitalized for either a heart attack, unstable angina or worsening symptoms of stable coronary artery disease. Their median age was 57 years, 81 percent were male, 40 percent had suffered a heart attack and 25 percent had diabetes.

Researchers visited the patients two to four weeks after their discharge, and then three more times at approximately three-month intervals. At each visit, a portable electrocardiograph called a Holter monitor recorded the patients' heart activity for 24 hours. All participants were



monitored on the first visit, and 35 had monitoring on more than one visit. Researchers averaged monitor readings over each half-hour, providing 5,979 half-hour observations. They then examined the relation of these ECG measurements with levels of several pollutants, including black carbon, produced by the incomplete combustion of fossil fuels, and particulate matter (PM2.5) of less than 2.5 micrometers (about 1/10,000th of an inch) in diameter.

Researchers obtained the PM2.5 and black carbon readings at a Harvard School of Public Health monitoring site, an average distance of 10.9 miles from the participants' homes.

Among the study's findings:

-- Increased levels of particular pollution – PM2.5 and black carbon (a marker for traffic exhaust) – were associated with ST-segment depression in the study participants.

-- Sulfur dioxide, a pollutant that can have non-traffic sources, also was associated with ST-segment depression.

-- No significant correlation was found between ST-segment depression and increased levels of carbon monoxide, but levels of this pollutant were low in this study.

-- Patients recovering from a heart attack had greater changes in ST segment depression on electrocardiograms compared to other participants.

The key question remains – how breathing air polluted by PM2.5 and black carbon might cause ST segment depression.

"Further research is needed to evaluate whether the pollution-related STsegment depression that we see is related to increased heart muscle inflammation, reduced oxygen flow, oxidative stress, or increased risk of arrhythmias," Gold said.



"We think that our findings, which are definitely subclinical, may represent a process that increases clinical risk for people with symptomatic coronary artery disease," she said.

Source: American Heart Association

Citation: Air pollution can hinder heart's electrical functioning (2008, September 8) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2008-09-air-pollution-hinder-heart-</u> <u>electrical.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.