

How to differentiate benign from malignant bile duct strictures?

September 9 2008

The differentiation of benign and malignant strictures is difficult. Recently, a group of clinical specialists in Netherlands attempted to find possible criteria for differentiation of malignant from benign bile duct strictures. They found that except for vascular involvement which was associated significantly with malignancy, there were no conclusive features of malignancy on regular imaging modalities.

The main etiology of bile duct strictures closely related to the liver is a malignancy (cancer). However, the differentiation of benign and malignant strictures is notoriously difficult. The consequences for the patient are considerable because cancer in this anatomical location requires extensive surgery with removal of a large part of the liver.

Extensive work-up including multi-slice computed tomography (CT), colour Doppler ultrasonography and magnetic resonance imaging (MRI) may improve the diagnostic dilemma. However, up to date, no single investigation reliably differentiates malignant from benign bile duct strictures.

A research article to be published on August 28, 2008 in the *World Journal of Gastroenterology* addresses this question. The research team led by Prof. van Gulik from the Surgical Department, analyzed a patient cohort that underwent resection for presumed malignancy of the bile duct, during the period 1998 to 2006. The final histologic diagnosis of 68 patients was correlated with the preoperative clinical, laboratory and radiological findings. Fifteen percent patients were found to have a

benign lesion. T

he findings of the different imaging studies revealed that only one feature, i.e. involvement of the blood vessels, showed a significant association with malignancy. All other features including clinical presentation, laboratory tests, brush cytology, and other imaging studies were unable to differentiate malignant and benign strictures.

Despite using recent, state-of-the-art imaging modalities, 15% patients with presumed malignancy underwent an extensive operation but were ultimately diagnosed to have benign strictures. Due to the relatively low incidence of vascular involvement (31% in this series), the absence of this finding is not conclusive. The article demonstrates that the differentiation of bile duct strictures remains difficult and that more research is needed to solve this diagnostic problem.

Source: World Journal of Gastroenterology

Citation: How to differentiate benign from malignant bile duct strictures? (2008, September 9) retrieved 3 May 2024 from

<https://medicalxpress.com/news/2008-09-differentiate-benign-malignant-bile-duct.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
