

## New classification of spinal deformity defines range of normalcy

December 9 2008

A University of Cincinnati (UC) neurosurgeon who has spent his career helping people with severe spine problems stand up straight has spearheaded the creation of a new spinal deformity classification system. The system, published this fall in the journal *Neurosurgery*, defines deformity in relation to the healthy, normal curve of the spine.

"What we've done is define spinal deformity and its manifestations throughout the course of a lifetime, based on a systematic approach to the spine, from the head to the pelvis," says Charles Kuntz IV, MD, an associate professor in UC's neurosurgery department and director of the division of spine and peripheral nerve surgery at the UC Neuroscience Institute. "Defining deformity with this degree of precision allows us to provide optimal treatment."

Kuntz, who practices at the Mayfield Clinic, and his co-authors defined spinal deformity by synthesizing published literature that describes normal neutral upright spinal alignment in asymptomatic juvenile, adolescent, adult and geriatric volunteers. The researchers used a total of 38 angles and displacements to define neutral upright spinal alignment, compiling their data over a period of five years.

The spine is a "dynamic organ that changes during the course of a lifetime," Kuntz says, with normal curves increasing with age.

An estimated 1.5 percent of the population has some degree of spinal deformity, which can take many forms. Abnormal curvatures can occur



from side to side, as in scoliosis; they can involve an abnormal forward curve of the spine, known as kyphosis, or hunchback; and they can involve an abnormal posterior curve of the lower spine, known as lordosis, or swayback.

Spinal deformity, depending on its severity, can cause pain, disability and a reduction in quality of life.

Kuntz, whose spinal reconstructions can take as long as 10 to 15 hours over a period of two days, strives for optimal spinal alignment with the finest cosmetic symmetry, even in the most severely disabled patients.

"Some physicians may feel that the result doesn't have to be perfect," Kuntz says. "But I do. It's a big deal when you have a patient who can't stand up straight, who can't look you in the eye, who's embarrassed to go out. It's a big deal when you help him or her become a person who's not only attractive to others but also attractive to himself or herself."

Source: University of Cincinnati

Citation: New classification of spinal deformity defines range of normalcy (2008, December 9) retrieved 13 May 2024 from <u>https://medicalxpress.com/news/2008-12-classification-spinal-deformity-range-normalcy.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.