

# MRI shows new types of injuries in young gymnasts

December 1 2008

---

Adolescent gymnasts are developing a wide variety of arm, wrist and hand injuries that are beyond the scope of previously described gymnastic-related trauma, according to a study presented today at the annual meeting of the Radiological Society of North America (RSNA).

"The broad constellation of recent injuries is unusual and might point to something new going on in gymnastics training that is affecting young athletes in different ways," said the study's lead author, Jerry Dwek, M.D., an assistant clinical professor of radiology at the University of California, San Diego and a partner of San Diego Imaging at Rady Children's Hospital and Health Center.

Previous studies have reported on numerous injuries to the growing portion of adolescent gymnasts' bones. However, this study uncovered some injuries to the bones in the wrists and knuckles that have not been previously described. In addition, the researchers noted that these gymnasts had necrosis, or "early death," of the bones of their knuckles.

"These young athletes are putting an enormous amount of stress on their joints and possibly ruining them for the future," Dr. Dwek said.

The radius is the bone in the forearm that takes the most stress during gymnastics. Due to damage to the radial growth plates, the bone does not grow in proportion to the rest of the skeleton and may be deformed. Consequently, it is not unusual for gymnasts to have a longer ulna than radius. Some former gymnasts must undergo surgery to shorten the ulna

and regain the proper fit of the wrist bones into the forearm.

Dr. Dwek and coauthor Christine Chung, M.D., used MRI to study overuse injuries seen in the skeletally immature wrists and hands of gymnasts. The researchers studied wrist and hand images of 125 patients, age 12 to 16, including 12 gymnasts with chronic wrist or hand pain.

"We were surprised to be looking at injuries every step down the hand all the way from the radius to the small bones in the wrist and on to the ends of the finger bones at the knuckles," Dr. Dwek said. "These types of injuries are likely to develop into early osteoarthritis."

Dr. Dwek suggested that additional study is needed to understand how gymnastic stresses are causing these injuries.

"It is possible that by changing the way that practice routines are performed, we might be able to limit the stress on the joints and on delicate growing bones," he said.

Source: Radiological Society of North America

Citation: MRI shows new types of injuries in young gymnasts (2008, December 1) retrieved 19 April 2024 from <https://medicalxpress.com/news/2008-12-mri-injuries-young-gymnasts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.