

# Have numbness, pain or muscle weakness? Guidelines identify best tests for neuropathy

December 3 2008

---

New guidelines developed by the American Academy of Neurology find a combination of blood tests and other specialized assessments appear to be the most helpful tests for finding the cause of neuropathy. Also known as neuritis or distal symmetric polyneuropathy, this common nerve problem affects people of all ages. The guidelines are published in the December 3, 2008, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

Neuropathy affects one in 50 people in the general population and one in 12 over the age of 55. It usually causes numbness, tingling or pain, often starting in the feet and moving to the hands. Symptoms spread slowly and evenly up the legs and arms. Muscle wasting and weakness can also occur. Neuropathy takes many forms and has many causes. The most common cause is diabetes. Other common causes are heredity, alcohol abuse, poor nutrition and autoimmune processes. Not all of the causes are known.

"There are many people with a neuropathy who have been walking around for years without having been diagnosed and treated," said guidelines author John D. England, MD, Chairman of the Department of Neurology at Louisiana State University Health Sciences Center in New Orleans and Fellow of the American Academy of Neurology. "Both neurologists and people with neuropathy need to know that the appropriate choice of tests is critical to accurate diagnosis."

To develop the guidelines, the authors analyzed all available scientific

studies on the topic.

The guidelines recommend that doctors obtain certain blood tests for all patients with numb, painful feet. "People with suspected nerve problems should talk to their doctors about screening tests, especially blood glucose, vitamin B12 level and serum protein levels, since these tests can often point to common causes of neuropathy," said England. The guidelines recommend tailored genetic testing for accurately diagnosing certain neuropathies that run in families.

The guidelines further recommend that doctors consider a combination of specialized tests to accurately evaluate neuropathies with autonomic dysfunction. These tests, known as autonomic tests, measure the action of the tiny nerves that control such functions as sweating, heart rate and blood pressure. Skin biopsy may also be useful to diagnose loss of tiny nerve fibers in the skin.

Source: American Academy of Neurology

Citation: Have numbness, pain or muscle weakness? Guidelines identify best tests for neuropathy (2008, December 3) retrieved 25 April 2024 from <https://medicalxpress.com/news/2008-12-numbness-pain-muscle-weakness-guidelines.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.