

Overweight siblings of children with type 2 diabetes likely to have abnormal blood sugar levels

December 9 2008

Overweight siblings of children with type 2 diabetes are four times more likely to have abnormal glucose levels compared to other overweight children. Because abnormal glucose levels may indicate risk for diabetes or diabetes itself, these children could benefit from screening tests and diabetes prevention education.

Researchers from The Children's Hospital of Philadelphia published their findings today in the online edition of the Journal of Pediatrics.

"To our knowledge, previous studies have not specifically looked at the risk of abnormal glucose tolerance among siblings of children diagnosed with type 2 diabetes. This group has a unique combination of genetic and environmental risk factors," said Sheela N. Magge, M.D., M.S.C.E., a pediatric endocrinologist at The Children's Hospital of Philadelphia and primary author of the study. "Clinical experience suggests that children with type 2 diabetes often have an obese sibling, which makes siblings an appropriate target for prevention trials."

The study looked at 62 children: 20 obese subjects with a sibling who had type 2 diabetes and a control group of 42 obese children. The groups were similar for age, gender, racial distribution (predominantly African American), pubertal status and body mass index over 95th percentile.

The researchers found that overweight siblings of children with type 2

diabetes had four times greater odds of having abnormal glucose levels (impaired glucose tolerance or type 2 diabetes) than other overweight children. However, investigators found no significant differences in insulin resistance, as measured by the homeostasis model assessment.

Type 2 diabetes is caused by a combination of both genetic and environmental factors. Known risks include obesity, decreased physical activity, race/ethnicity, family history and insulin resistance. Obesity decreases insulin sensitivity, as does puberty, when all adolescents experience a period of relative insulin resistance. In obese adolescents already at risk of developing type 2 diabetes, the increase in insulin resistance during puberty may be enough to unmask disease. Family history is also important; 74 to 100 percent of children with type 2 diabetes have a first- or second-degree relative who also has the condition.

Not all children with a family history of type 2 diabetes, insulin resistance or obesity develop type 2 diabetes, cautions Dr. Magge.

The researchers also add that identifying groups at high risk for type 2 diabetes during childhood, such as obese siblings of children with type 2 diabetes, could help guide screening of obese children for abnormal glucose tolerance by primary care providers. This could also help to identify children who might benefit from participation in future type 2 diabetes prevention studies.

Source: Children's Hospital of Philadelphia

Citation: Overweight siblings of children with type 2 diabetes likely to have abnormal blood sugar levels (2008, December 9) retrieved 16 July 2024 from <https://medicalxpress.com/news/2008-12-overweight-siblings-children-diabetes-abnormal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.