

Study finds genetic link between sleep disorders and depression in young children

February 1 2009

A study in the Feb. 1 issue of the journal *Sleep* was the first to use twin data to examine the longitudinal link between sleep problems and depression. Results of this study demonstrate that sleep problems predict later depression; the converse association was not found. These findings are consistent with the theory that early treatment of sleep problems may protect children from the development of depression.

Results of the study indicate that the stability of sleep problems across time is mainly caused by genetic factors (46 percent of the genetic influences on sleep at age 10 were the same as those that influenced a child at age 8). The stability of depression is mainly caused by non-shared environmental influences (19 percent of the non-shared environmental influence on depression at age 10 remained the same from the age of eight).

According to lead author Alice M. Gregory, senior lecturer in the department of psychology at Goldsmiths College in London, the most surprising result of the study concerned the reasons why there may be links between sleep problems and depression at different points in a young person's life.

"We reported in a study previously, that genes were the most important factor in explaining the association between sleep problems and depression in eight year olds," said Gregory. "However, when we examined this issue at age 10, we found that genes were less important in explaining the association and that environmental influences had become

more important. This could be because environmental experiences are becoming more relevant as children grow older and could therefore play a role in both sleep problems and depression."

The process thought to underlie the associations between sleep and depression reflects abnormalities that are known to be influenced by genetic variations. A longitudinal twin study was used in order to distinguish shared and non-shared environmental influences. Shared environmental influences create similarities among individual family members, and can therefore be estimated by calculating the similarity between identical twins that is not due to genetic influences. Non-shared environmental influences individualize family members, and can therefore be estimated by examining the associations between monozygotic (MZ) twin pairs; if there is not a perfect correlation between MZ twins, it can be assumed that non-shared environmental influence are playing a role.

Data were collected from 300 twin pairs born in England and Wales between 1994 and 1996. Anxiety was examined at age 7 using the parent-report Anxiety Related Behaviors Questionnaire (ARBQ), a 21-item scale reflecting commonly assessed anxiety-related behaviors in young children (including general distress, separation anxiety, shyness, inhibition and fears). Two hundred forty-seven twin pairs scored highly on parent-reported anxiety at age seven; fifty-three control twin pairs also were selected to ensure full coverage of the full range of scores on test measures.

Parents reported on their 8-year-old children's sleep problems using the abbreviated version of the Child Sleep Habits Questionnaire (CSHQ). Eight aspects of commonly reported childhood sleep problems were analyzed, including bedtime resistance, sleep-onset delay, sleep duration, sleep anxiety, night waking, parasomnias, sleep-disordered breathing and daytime sleepiness.

Depressive symptoms were examined in the 8-year-old twins through the Children's Depression Inventory (CDI), a 27-item self-report questionnaire that examines affective, cognitive and behavioral signs of depression. Approximately two years later when the twins were 10 years old, data on sleep problems and depression were collected again from 250 of the twin pairs.

The authors argue that although childhood sleep problems have only a small influence on later depression, these issues should be given serious consideration as they can negatively affect a child's mood, attention-span, and social and academic function. They also state that, in comparison with other risk indicators of later problems, sleep difficulties are more easily broached and readily discussed with families without the negative stigma that may be associated with the discussion of mental health problems in children.

Source: American Academy of Sleep Medicine

Citation: Study finds genetic link between sleep disorders and depression in young children (2009, February 1) retrieved 2 May 2024 from <https://medicalxpress.com/news/2009-02-genetic-link-disorders-depression-young.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--