

MRI shows brain atrophy pattern that predicts Alzheimer's

February 10 2009

Using special MRI methods, researchers have identified a pattern of regional brain atrophy in patients with mild cognitive impairment (MCI) that indicates a greater likelihood of progression to Alzheimer's disease. The findings are published in the online edition of *Radiology*.

"Previously, this pattern has been observed only after a diagnosis of probable Alzheimer's disease," said the study's lead author, Linda K. McEvoy, Ph.D., assistant project scientist in the Department of Radiology at the University of California San Diego School of Medicine in La Jolla. "Our results show that some individuals with MCI have the atrophy pattern characteristic of mild Alzheimer's disease, and these people are at higher risk of experiencing a faster rate of brain degeneration and a faster decline to dementia than individuals with MCI who do not show that atrophy pattern."

According to the Alzheimer's Association, more than five million Americans currently have Alzheimer's disease. One of the goals of modern neuroimaging is to help in early and accurate diagnosis, which can be challenging. There is no cure for Alzheimer's disease, but when it is diagnosed early, drug treatment may help improve or stabilize patient symptoms.

In Alzheimer's disease, nerve cell death and tissue loss cause areas of the brain to atrophy. Structural MRI allows radiologists to visualize subtle anatomic changes in the brain that signal atrophy. MCI is associated with an increased risk of progression to Alzheimer's disease. Rates of

progression vary. Some patients progress rapidly, while others remain stable for relatively long periods of time.

For the study, Dr. McEvoy and colleagues set out to determine if they could identify a pattern of regional atrophy characteristic of mild Alzheimer's disease in order to aid in the prediction of cognitive decline in patients with MCI.

In the study, the researchers analyzed brain MR images from 84 patients with mild Alzheimer's disease, 175 patients with MCI and 139 healthy controls, using with semi-automated, individually specific quantitative MRI methods. The results showed widespread cortical atrophy in some patients with MCI, involving all cortical areas except those involved with processing of primary motor and sensory information. However, most indicative of future cognitive decline were atrophy in parts of the medial and lateral temporal lobes and in the frontal lobes. This pattern was also present in the patients with mild Alzheimer's disease.

"Although these individuals are reporting problems mainly with memory, the atrophy involves more than just memory areas, extending into brain regions involved in planning, organization, problem solving and language," Dr. McEvoy said.

Follow-up data were available for 160 patients with MCI. The patients exhibiting atrophy in the brain regions described above showed significant one-year clinical decline and structural brain loss and were more likely to progress to a probable diagnosis of Alzheimer's disease. MCI patients without that pattern of atrophy remained stable after a year.

Dr. McEvoy hopes that these findings will have an impact on the design of clinical trials to test medications that may slow or halt the progression of Alzheimer's disease.

"Currently there are no treatments that will prevent or cure Alzheimer's disease, but information about risk of rapid decline may help patients with MCI and their families plan for the future," Dr. McEvoy said.

Source: Radiological Society of North America

Citation: MRI shows brain atrophy pattern that predicts Alzheimer's (2009, February 10)
retrieved 27 April 2024 from
<https://medicalxpress.com/news/2009-02-mri-brain-atrophy-pattern-alzheimer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.