

Time of conception linked to birth defects in United States

March 30 2009

A study published in the April 2009 issue of the medical journal *Acta Paediatrica* is the first to report that birth defect rates in the United States were highest for women conceiving in the spring and summer. The researchers also found that this period of increase risk correlated with increased levels of pesticides in surface water across the United States.

Studying all 30.1 million births which occurred in the U.S. between 1996 and 2002, the researchers found a strong association between the increased number of [birth defects](#) in children of women whose last menstrual period occurred in April, May, June or July and elevated levels of nitrates, atrazine and other pesticides in [surface water](#) during the same months. While many of these chemicals, including the herbicide atrazine which is banned in European countries but permitted in the U.S., are suspected to be harmful to the developing embryo, this is the first study to link their increased seasonal concentration in surface water with the peak in birth defects in infants conceived in the same months.

The correlation between the month of last menstrual period and higher rates of birth defects was statistically significant for half of the 22 categories of birth defects reported in a Centers for Disease Control database from 1996 to 2002 including spina bifida, [cleft lip](#), clubfoot and Down's syndrome.

"Elevated concentrations of pesticides and other agrochemicals in surface water during April through July coincided with significantly

higher risk of birth defects in live births conceived by women whose last menstrual period began in the same months. While our study didn't prove a cause and effect link, the fact that birth defects and pesticides in surface water peak during the same four months makes us suspect that the two are related," said Paul Winchester, M.D., Indiana University School of Medicine professor of clinical pediatrics, the first author of the study.

"Birth defects, which affect about 3 out of 100 newborns in the U.S., are one of the leading causes of infant death. What we are most excited about is that if our suspicions are right and pesticides are contributing to birth defect risk, we can reverse or modify the factors that are causing these lifelong and often very serious medical problems," said Dr. Winchester, a Riley Hospital for Children neonatologist.

Birth defects are known to be associated with risk factors such as alcohol, smoking, diabetes or advanced age. However, the researchers found that even mothers who didn't report these risk factors had higher overall birth defect rates for babies conceived from April to July.

The study relies on findings by U.S. Geological Survey, the U.S. Environmental Protection Agency and other agencies on the seasonal variations in nitrates, atrazine and other pesticides in the surface water.

"These observations by Dr. Winchester are extremely important, as they raise the question for the first time regarding the potential adverse effect of these commonly used chemicals on pregnancy outcome - the health and well-being of our children," said James Lemons, M.D., Hugh McK. Landon Professor of Pediatrics at the IU School of Medicine. Dr. Lemons is director of the section of neonatal-perinatal medicine at Riley Hospital.

Source: Indiana University ([news](#) : [web](#))

Citation: Time of conception linked to birth defects in United States (2009, March 30) retrieved 2 May 2024 from

<https://medicalxpress.com/news/2009-03-conception-linked-birth-defects-states.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.