

# Before starting dialysis, patients need nephrologist care

March 25 2009

---

For patients with end-stage renal disease (ESRD), receiving care from a nephrologist in the months before starting dialysis reduces the risk of death during the first year on dialysis, reports a study in the May 2009 issue of the *Journal of the American Society of Nephrology* (JASN). The study also shows geographic "clusters" where pre-dialysis care for patients with advanced chronic kidney disease (CKD) is not optimal. "Assistance to improve pre-dialysis care might be profitably targeted to specific treatment centers and the health care systems they serve," comments William McClellan, MD (Emory University School of Medicine, Atlanta, GA).

Dr. McClellan and colleagues analyzed data on more than 30,000 [patients](#) starting [dialysis](#) in five of the 18 US ESRD Network regions. The researchers evaluated the quality of the patients' medical care in the months before their CKD progressed to ESRD, and how that affected the patients' outcomes on dialysis.

Just over half of the patients received at least six months of pre-dialysis care from a nephrologist, as recommended by current guidelines. For these patients, the chances of surviving the first year on dialysis were about 50 percent higher than for patients who did not receive at least six months of nephrologist care. Survival rates were higher at dialysis centers where more patients received recommended care.

The researchers also unexpectedly discovered that dialysis centers with the lowest rates of recommended pre-dialysis care tended to be

"clustered geographically." For example, there was a "significant circular cluster" of low pre-dialysis care centers located in Alabama and Mississippi.

Although the reasons for the geographic variations in care are unclear, the results identify specific regions that might benefit from efforts to improve care for advanced CKD patients. "The Centers for Medicare & Medicaid Services are currently conducting a pilot quality improvement initiative in ten states to determine the feasibility of such efforts," says Dr. McClellan.

More information: The study, entitled "Treatment Center and Geographic Variability in Pre-ESRD Care Associate with Increased Mortality," will appear online at [jasn.asnjournals.org/](http://jasn.asnjournals.org/) on Wednesday, March 25, 2009, doi 10.1681/ASN.2008060624.

Source: American Society of Nephrology ([news](#) : [web](#))

Citation: Before starting dialysis, patients need nephrologist care (2009, March 25) retrieved 3 May 2024 from <https://medicalxpress.com/news/2009-03-dialysis-patients-nephrologist.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--