


Evidence appears to show how and where frontal lobe works

March 2 2009



The neuroscience of decision-making By examining stroke victims, researchers determined that the brain's frontal lobe controls decision-making, with abstract decisions made closer to the front and concrete decisions farther back. Image: Journal of Cognitive Neuroscience

(Physorg.com) -- A Brown University study of stroke victims has produced evidence that the frontal lobe of the human brain controls decision-making along a continuum from abstract to concrete, from front to back.

Abstract actions can be controlled at an abstract level, such as deciding to make a sandwich, or at more concrete and specific levels, such as choosing a sequence of movements that make the sandwich.

The scientific data supports preexisting theories that abstract decisions about action take place in the front of the frontal lobe, the back portion controls the capacity for concrete decisions, and the progression from front to back forms a gradient from abstract to concrete.

The Brown researchers are among the first to show that specific areas of the frontal cortex are needed for different levels of abstract decision.

The finding, to be detailed March 1 in the journal *Nature Neuroscience*, represents a huge leap in comprehending how the brain supports higher level cognition and intelligent behavior. It could lead to advances in everything from the treatment of strokes to understanding how humans develop thought. Researchers from the University of California-Berkeley also participated in the study.

"It is among the strongest evidence to date for a systemic organization of the frontal cortex," said lead author David Badre, an assistant professor of cognitive and linguistic sciences at Brown University.

The frontal cortex of brain has been long known to affect the internal control of behavior. It controls the capacity to plan, reason, conduct higher-level thinking and connect what we know about the world to how we behave.

Badre and his collaborators came to their conclusion by studying stroke victims who suffered damage to different parts of the frontal lobe. The patients all suffered a stroke at least six months prior to testing. All were screened with an MRI or CT scan to determine where any lesions existed in the brain post-stroke.

The scientists recruited 11 patients — seven men and four women, ranging from age 45 to 73. A 12th patient was recruited but could not perform any of the tests involved.

Researchers gave the patients four different tests that ultimately required selecting a finger-press response. For example, the first test would show a color such as red, which required an index finger push. Blue would trigger the middle finger. The test would then become more difficult by adding more alternate finger presses.

Patients faced greater challenges in selecting a response as subsequent, progressive tests became more complex, with more abstract options.

Badre and colleagues found that damage at a given location affected more abstract decisions but left intact the capacity for more concrete decisions. "If there is damage in a given spot, it will affect all higher (decision-making) functions but not lower functions," Badre said.

Provided by Brown University

Citation: Evidence appears to show how and where frontal lobe works (2009, March 2) retrieved 27 April 2024 from <https://medicalxpress.com/news/2009-03-evidence-frontal-lobe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--