

Researchers find link between anesthesia exposure and learning disabilities in children

March 24 2009

Mayo Clinic researchers have found that children who require multiple surgeries under anesthesia during their first three years of life are at higher risk of developing learning disabilities later. Several studies have suggested that anesthetic drugs may cause abnormalities in the brains of young animals. This is the first study in humans to suggest that exposure of children to anesthesia may have similar consequences. The finding is reported in the current issue of the journal [Anesthesiology](#).

Using data from the long-term [Rochester Epidemiology Project](#), researchers studied the medical records of 5,357 [children](#) from Olmsted County who were born between 1976 and 1982.

The research team, led by Robert Wilder, M.D., Ph.D., a [Mayo Clinic](#) anesthesiologist, found that although one exposure to [anesthesia](#) was not harmful, more than one almost doubled the risk that a child would be identified as having a [learning disability](#) before age 19. The risk also increased with longer durations of anesthesia.

"It's very important for parents and families to understand that although we see a clear difference in the frequency of learning disabilities in children exposed to anesthesia, we don't know whether these differences are actually caused by anesthesia," says Randall Flick, M.D., a Mayo Clinic anesthesiologist and co-author of the study.

"The problem is that anyone who underwent an anesthetic also had surgery," says Dr. Wilder. "It's unclear whether it's the anesthetic, the

physiological stress of surgery or perhaps the medical problems that made surgery necessary that are responsible for the learning disabilities."

Young children's brains are more vulnerable to a variety of problems because they are undergoing dynamic growth. The brain is rapidly forming connections between cells and trimming excess cells and connections, says Dr. Wilder.

The general anesthesia chemicals in use during the study period were primarily halothane and [nitrous oxide \(laughing gas\)](#). Although halothane is no longer used in the U.S., it has been replaced by newer agents that have similar effects on the brain. Nitrous oxide is widely used throughout the U.S. and the world.

Debate exists about the developmental correlation between the animal (rodent) and human studies. Some think that the related exposure period would be perinatal in humans (the last month of pregnancy and first six months after birth), so the researchers repeated their analysis, examining anesthetic exposure before age 2, and found similar results.

"Parents and physicians need to balance this information along with the normal decisions that we all go through when we decide to have surgery for one of our children," says Dr. Flick. "Although alternatives to the use of these medications exist, they are limited. Certainly, performing surgery without appropriate use of anesthesia is unacceptable."

The children in the study were tested as a natural part of the educational process in the Rochester school system. They did not perform as well in reading, writing or math as their IQ tests indicated.

Other studies have linked anesthesia exposure in young children to behavioral problems. Dr. Flick says the Food and Drug Administration (FDA) is aware of the possible problems with anesthesia. "They've been

very proactive in trying to gather information as quickly and thoughtfully as possible," Dr. Flick says, "but much more research is needed before we could conclude that anesthesia itself causes problems." He also encourages families with questions to go to the Web sites of the American Society of Anesthesiology and the Society for Pediatric Anesthesia.

The research team is working to obtain funding to extend the database for 10 more years (1982-1992), a period that would include the use of more modern anesthetics. They are also working with the FDA to complete a study that matches children who had an anesthetic with children who have a similar medical problem but did not receive an anesthetic.

Source: Mayo Clinic ([news](#) : [web](#))

Citation: Researchers find link between anesthesia exposure and learning disabilities in children (2009, March 24) retrieved 10 April 2024 from <https://medicalxpress.com/news/2009-03-link-anesthesia-exposure-disabilities-children.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--