

## Antioxidant found in berries, other foods prevents UV skin damage that leads to wrinkles

April 21 2009

---

Using a topical application of the antioxidant ellagic acid, researchers at Hallym University in the Republic of Korea markedly prevented collagen destruction and inflammatory response - major causes of wrinkles -- in both human skin cells and the sensitive skin of hairless mice following continuing exposure to UV-B, the sun's skin-damaging ultraviolet radioactive rays.

Ji-Young Bae, a graduate student in the laboratory of Dr. Young-Hee Kang, presented results of the two-part study on Tuesday, April 21, at the Experimental Biology 2009 meeting in New Orleans. The presentation was part of the scientific program of the American Society for Nutrition.

Ellagic [acid](#) is an antioxidant found in numerous fruits, vegetables and nuts, especially raspberries, strawberries, cranberries and pomegranates. Earlier studies have suggested it has a photoprotective effect.

But how? The Kang laboratory found that, in human skin cells, ellagic acid worked to protect against UV damage by blocking production of MMP (matrix metalloproteinase enzymes that break down collagen in damaged [skin cells](#)) and by reducing the expression of ICAM (a molecule involved in inflammation).

The scientists then turned to young (four weeks), male, hairless mice -

genetically bred types of mice often used in dermatology studies because of the physiological similarities of their skin to that of humans. For eight weeks, the 12 mice were exposed to increasing [ultraviolet radiation](#), such as that found in sunlight, three times a week, beginning at a level sufficient to cause redness or sunburn and increasing to a level that would have definitely caused minor skin damage to [human skin](#).

During these eight weeks, half of the exposed mice were given daily 10 microM topical applications of ellagic acid on their skin surface, even on the days in which they did not receive UV exposure. The other mice, also exposed to UV light, did not receive ellagic acid. (Another six mice served as controls, with neither UV exposure nor ellagic acid.)

What happened? First, as expected, the mice exposed to UV radiation without the ellagic acid treatment developed wrinkles and thickening of the [skin](#).

Second, as hypothesized, the exposed [mice](#) that received topical application of ellagic acid showed reduced wrinkle formation.

Third, as suggested in the study of human cells, the ellagic acid reduced inflammatory response and MMP secretion due to protection from the degradation of collagen. The ellagic acid also helped prevent an increase of epidermal thickness

The researchers say the results demonstrate that ellagic acid works to prevent wrinkle formation and photo-aging caused by UV destruction of collagen and [inflammatory response](#).

Source: Federation of American Societies for Experimental Biology  
([news](#) : [web](#))

Citation: Antioxidant found in berries, other foods prevents UV skin damage that leads to wrinkles (2009, April 21) retrieved 20 April 2024 from <https://medicalxpress.com/news/2009-04-antioxidant-berries-foods-uv-skin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.