

## Gene test determines risk of heart surgery complications

## April 30 2009

Genetic differences can explain why some patients undergoing heart surgery later experience shock and kidney complications, according to a study by researchers at the Charité - Universitätsmedizin Berlin, the Max-Delbrück-Center for Molecular Medicine (MDC) Berlin-Buch in Germany and the Austin Hospital in Melbourne, Australia. The results indicate that performing a genetic test on patients before they have surgery can help guide treatment after they leave the operating room.

The researchers studied the gene that encodes the <u>enzyme</u> catechol-Omethyltransferase (COMT). Certain variants of the COMT gene have long been suspected to play a role in shock and <u>kidney failure</u> in patients following <u>heart surgery</u>. The COMT enzyme is involved in metabolizing norepinephrine (noradrenalin), a drug that is given to patients postsurgery to stimulate their blood flow and to normalize their blood pressure.

Professor Duska Dragun, MD, Charité, Professor Friedrich Luft, MD (Experimental and Clinical Research Center, MDC) and Dr. Wolf-Hagen Schunck (MDC) studied the COMT gene in 260 patients who underwent heart bypass surgery. They were able to show that the genetic variant they call "LL" can lower the activity of the COMT enzyme. As a result, LL patients are more likely to develop shock and kidney failure.

In addition, LL patients who experience shock do not respond very well to treatment with norepinephrine. Since the activity of the COMT enzyme is lowered in LL patients, norepinephrine cannot fully be



metabolized. As a result, too much norepinephrine remains in the body and the drug is no longer effective.

Therefore, the researchers suggest that "perhaps, more suitable hemodynamics could be achieved in LL patients were they given vasopressin rather than noradrenaline (norepinephrine) and acute kidney injury might be attenuated by avoidance of cardiopulmonary bypass and nephrotoxic medication." Larger clinical trials are necessary to show whether patients need to undergo genetic testing prior to heart surgery to determine their risk for shock and kidney failure.

More information: Journal of the American Society of Nephrology, JASN, doi 10.1681/ASN.2008080915

Source: Helmholtz Association of German Research Centres (<u>news</u> : <u>web</u>)

Citation: Gene test determines risk of heart surgery complications (2009, April 30) retrieved 30 April 2024 from https://medicalxpress.com/news/2009-04-gene-heart-surgery-complications.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.