

## Heart screening unnecessary in type 2 diabetes patients with no symptoms

April 14 2009

---

Routine screening for coronary artery disease in type 2 diabetes patients with no symptoms of angina or a history of coronary disease is unnecessary and may lead initially to more invasive and costly heart procedures, according to researchers at Yale School of Medicine. They report their findings in the April 15 *Journal of the American Medical Association*.

"Patients with type 2 diabetes with no symptoms who are feeling well can generally be managed effectively with preventive therapies such as lipid-lowering drugs, [blood pressure](#) medication, aspirin and diabetes treatment," said Lawrence Young, M.D., professor of medicine at Yale School of Medicine.

[Coronary artery disease](#) is a major cause of death and disability in patients with type 2 diabetes, who often undergo routine screening with stress testing. In this study, Young, Frans J. Wackers, M.D., Silvio Inzucchi, M.D., Deborah Chyun, and other colleagues sought to determine whether routinely screening diabetic patients without a history of heart problems helped identify those at higher [cardiac risk](#).

The team spearheaded the Detection of Ischemia in Asymptomatic Diabetics study in which 1,123 participants were randomly assigned to either a screening [stress test](#) that looks at the blood flow to the heart, or to a group that received no screening. Participants enrolled at 14 centers in the United States and Canada and were evaluated over the next five years.

The team found that routine screening of type 2 diabetics identifies a small group of patients at higher risk, but does not significantly affect overall outcome. With each test costing about \$1,000, and with 200 million patients with diabetes worldwide, Young says, "Routine screening would have tremendous economic implications and our findings did not indicate that routine stress testing had additional benefit in this population." The study also found that routine screening leads initially to more invasive procedures such as cardiac catheterization, stents or bypass.

"On the other hand," he cautioned, "stress testing has an important role in evaluating patients with [type 2 diabetes](#) who have symptoms suggesting heart disease, since these individuals are at very high risk."

"This is a bit of good news in the field of diabetes where patients and their physicians have real concern for heart disease. In patients getting modern medical therapy in our study, serious heart problems were infrequent," said Young.

More information: JAMA 301[15]:1547-1555 (April 15, 2009)

Source: Yale University ([news](#) : [web](#))

Citation: Heart screening unnecessary in type 2 diabetes patients with no symptoms (2009, April 14) retrieved 19 April 2024 from <https://medicalxpress.com/news/2009-04-heart-screening-unnecessary-diabetes-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.