

Study suggests that trouble sleeping leads to increased ratings of pain in cancer patients

April 15 2009

A study in the April 15 issue of the *Journal of Clinical Sleep Medicine* suggests that sleep problems lead to increased pain and fatigue in cancer patients. The results indicate that interventions aimed at trouble sleeping would be expected to improve both pain and fatigue in this patient population.

Results show that more than half the sample reported having trouble sleeping, with 26 percent reporting moderate or severe trouble sleeping. Compared with [patients](#) who reported no trouble sleeping, patients with moderate to severe trouble sleeping reported significantly more fatigue, pain and depressed mood. Using structural equation modeling analysis to evaluate causal relations and directions of effect, the best-fitting model indicates that trouble sleeping led to increased ratings of pain.

According to the authors, the relationship between pain and sleep often has been assumed to be reciprocal. In the present study, however, a model of reciprocal causation could not be fit to the data, and models in which pain caused trouble sleeping did not fit as well as the model in which trouble sleeping caused pain.

"We believed we would find a bi-directional relationship between [insomnia](#) and pain, but instead found that trouble sleeping was more likely a cause, rather than a consequence, of pain in patients with cancer," said lead author Edward J. Stepanski, chief operational officer at the Accelerated Community Oncology Research Network in Memphis, Tenn.

The study included demographic, clinical and patient-reported outcomes data from 11,445 [cancer patients](#) undergoing treatment at the West Clinic, a large community oncology practice in Memphis. Participants had an average age of 61.5 years, and 74 percent were female. [Breast cancer](#) was the most common form of cancer, and about 25 percent of study subjects had received [chemotherapy](#) in the last 30 days. Increases in depressed mood also led to increased ratings of pain.

Younger age and recent administration of chemotherapy were both associated with increased trouble sleeping. According to the authors, younger patients often receive more aggressive chemotherapy than older patients; therefore, younger patients may be exposed to more treatment-related toxicity.

Stepanski stated that several studies have shown that cognitive behavioral therapy (CBT) improves sleep in cancer patients who have insomnia. He believes that this type of intervention may decrease patients' [pain](#) and fatigue by improving their sleep.

More information: More information is available from the AASM about sleep and breast cancer at www.sleepeducation.com/Article.aspx?id=1076 and about cognitive behavioral therapy at www.sleepeducation.com/Treatment.aspx?id=5 .

Source: American Academy of Sleep Medicine ([news](#) : [web](#))

Citation: Study suggests that trouble sleeping leads to increased ratings of pain in cancer patients (2009, April 15) retrieved 25 April 2024 from <https://medicalxpress.com/news/2009-04-pain-cancer-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.