

# New ultrasound-guided biopsy method allows improved diagnosis of endometrial disease

April 14 2009

---



Dr. Elysia Moschos has shown that using saline-infusion sonography, or ultrasound, might make it easier to visualize and diagnose diseases in the lining of the uterus. Credit: UT Southwestern Medical Center

A procedure used in conjunction with a vaginal ultrasound might make it easier to visualize and diagnose diseases in the lining of the uterus, researchers at UT Southwestern Medical Center have found.

Physicians evaluated the endometrium, a cavity that lines the inside of the [uterus](#), in women who were in the midst of or had gone through menopause and who complained of abnormal bleeding. Abnormal bleeding can indicate certain diseases of the endometrium that may or

may not be malignant.

The current standard of care is to blindly biopsy the endometrium; however, the biopsy might not always sample the part of the cavity that is diseased.

A study available online and in the April issue of *Obstetrics and Gynecology* shows that using saline-infusion sonography (SIS), or ultrasound, a minimally invasive procedure, allows doctors to actually see where in the endometrium a polyp or growth exists and to biopsy it accordingly.

"Saline infusion sonography augments the usual transvaginal sonogram and lets us see what is inside the endometrium," said Dr. Elysia Moschos, associate professor of obstetrics and gynecology at UT Southwestern and the lead author of the study. "Normally, the endometrial cavity is collapsed, but when we infuse it with saline, we can visualize inside the cavity and see if it contains any abnormalities."

On a sonogram, water appears dark and tissue is light, so polyps or tumors look bright in comparison with the surrounding saline solution. Using the SIS procedure, physicians inject saline through a [catheter](#) threaded into the uterus through the cervix to fill up and expand the endometrial cavity.

Doctors can then easily visualize and biopsy an existing growth under sono guidance and send it to a pathologist for analysis.

A total of 88 saline-infusion sonography endometrial samples were obtained. In the final outcome of 80 of those samples, saline-infusion endometrial sampling provided a diagnosis 89 percent of the time, compared with 52 percent for endometrial biopsy.

The study showed there were no women for whom blind biopsy of the endometrium would still be an appropriate first step, Dr. Moschos said. The blind biopsies missed 15 of 16 benign polyps and one-third of cancers.

By comparison, two-thirds of benign [polyps](#) were correctly diagnosed by saline-infusion sonography endometrial sampling and no premalignant or malignant growths were missed.

Abnormal bleeding is a common complaint among patients, particularly during and after menopause. Women experiencing this symptom need to be evaluated for any kind of gynecologic cancers, Dr. Moschos said. The first step should be imaging of the endometrium by ultrasound.

According to the American College of [Obstetrics](#) and Gynecology, postmenopausal women who have an endometrial lining thicker than 4 millimeters need further follow-up.

"In women with abnormal endometrial appearances, SIS should then be performed. If the endometrium is uniformly thickened on the SIS exam, only then is a blind [biopsy](#) appropriate," Dr. Moschos said. "However, most endometrial pathology is focal, and therefore directed sampling, such as with saline infusion sonography guidance, is necessary."

Source: UT Southwestern Medical Center ([news](#) : [web](#))

Citation: New ultrasound-guided biopsy method allows improved diagnosis of endometrial disease (2009, April 14) retrieved 3 May 2024 from <https://medicalxpress.com/news/2009-04-ultrasound-guided-biopsy-method-diagnosis-endometrial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.