

Genes found to play a role in breast cancer's spread to the brain

May 6 2009

New research led by investigators at Memorial Sloan-Kettering Cancer Center (MSKCC) identifies three genes that specifically mediate the metastasis, or spread, of breast cancer to the brain and illuminates the mechanisms by which this spread occurs. The study was published online today in *Nature*.

According to the study, COX2 and HB-EGF — [genes](#) that induce cancer cell mobility and invasiveness — were found to be genetic mediators in the spread of [breast cancer](#) to the [brain](#). A third gene, ST6GALNAC5, was shown to provide cancer cells with the capability of exiting the [blood circulation](#) and passing through the blood-brain barrier to enter the brain tissue.

"Our research sheds light on the role these genes play in determining how breast tumor cells break free and, once mobile, how they decide where to attack," said Joan Massagué, PhD, Chair of the Cancer Biology and Genetics Program at MSKCC and a Howard Hughes Medical Institute investigator.

Breast cancer metastasis to the brain typically occurs years after removal of a breast tumor, suggesting that disseminated cancer cells initially lack the specialized functions needed to overtake the dense network of capillaries that constitute the blood-brain barrier. This barrier prevents the entry of circulating cells and regulates the transport of molecules into the brain tissue. To generate brain metastasis, circulating cancer cells must, therefore, be able to pass through the blood-brain barrier and

interact with the brain microenvironment.

In the study, Dr. Massagué and his colleagues isolated cancer cells that preferentially targeted the brain from patients with advanced disease. By combining this approach with gene expression profiling, additional testing in mouse model systems, and analysis of a body of clinical data, the investigators identified certain genes and functions that selectively mediate cancer cell passage through the blood-brain barrier.

The authors observed that ST6GALNAC5, an enzyme that is normally active only in [brain tissue](#), causes a chemical reaction that creates a coating on the surface of breast cancer cells that enhances their ability to breach the blood-brain barrier. Their findings show that breast cancer cells use this brain-specific cell-coating enzyme as a means of infiltrating the brain.

"Our results draw attention to the role of the cell surface coating as a previously unrecognized participant in brain metastasis, and to the possibility of using drugs to disrupt its interactions," said Dr. Massagué. "Further study is necessary to explore the role of these genes in brain metastasis and their interest as therapeutic targets."

The study authors also noted that COX2 and HB-EGF, which prime breast [cancer cells](#) for entrance into the brain, had previously been found to be involved in breast cancer's spread to the lung. This suggests a partial sharing of genetic mediators of metastasis to both the brain and lung and may explain the association of brain and lung relapse in women with breast cancer.

Source: Memorial Sloan-Kettering Cancer Center

Citation: Genes found to play a role in breast cancer's spread to the brain (2009, May 6) retrieved 2 May 2024 from <https://medicalxpress.com/news/2009-05-genes-role-breast-cancer-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.