

Study links ADHD with sleep problems in adolescents

May 1 2009

A study in the May 1 issue of the journal *SLEEP* shows that adolescents with a childhood diagnosis of Attention Deficit/Hyperactivity Disorder (ADHD) are more likely to have current and lifetime sleep problems and disorders, regardless of the severity of current ADHD symptoms. Authors suggest that findings indicate that mental health professionals should screen for sleep problems and psychiatric comorbidities among all adolescents with a childhood diagnosis of ADHD.

Results indicate that adolescents with a childhood diagnosis of ADHD, regardless of persistent ADHD were more likely to have current sleep problems and sleep disorders such as [insomnia](#), sleep terrors, nightmares, bruxism and snoring. Of the total sample, 17 percent of children with ADHD were currently suffering from primary insomnia, versus 7 percent of controls; lifetime primary insomnia occurred in 20 percent of children with ADHD, compared to 10 percent of controls. Nightmare disorder affected 11 percent of children with ADHD and lifetime nightmare disorder affected 23 percent, versus 5 and 16 percent of controls. The presence of at least one psychiatric comorbid condition increases the risks for insomnia and nightmares.

According to principal investigator Susan Shur-Fen Gau, MD, PhD, associate professor at the College of Medicine and Public Health, National Taiwan University, symptoms and consequences of ADHD and sleep problems in children often overlap. Some primary sleep disorders are found to be associated with inattention, hyperactivity, behavioral problems and impaired academic performance, which are often

mistaken for symptoms of ADHD.

"In some patients with ADHD, symptoms are caused or exaggerated by primary sleep disorders, and therefore treatment of the sleep disorder will improve ADHD symptoms," said Gau.

Data were collected from 281 consecutive patients (86.2 percent male) between the ages of 10 to 17 years who had been diagnosed with ADHD according to DSM-IV criteria at a mean age of 6.7 years, and 185 controls who did not have ADHD as a child or teen.

Diagnosis of ADHD was made based on information obtained from parent and child interviews, observation of the child's behaviors, and rating scales reported by parents and teachers.

Findings of the study indicated that the rates of nightmare and lifetime nightmare disorder were more prevalent in girls and snoring was more prevalent in boys. Snoring may be more prevalent in boys due to an increased rate of sleep-disordered breathing in boys. Mothers were found to be more aware of symptoms related to ADHD in the presence of primary insomnia, sleep terror disorder or sleepwalking disorder, whereas teachers may be more sensitive to ADHD symptoms in the presence of primary hypersomnia and nightmare disorder.

According to the study, sleep problems in children with ADHD may be caused by a variety of factors, including internet addiction, hyperactivity, use of stimulants and the presence of other psychiatric disorders. Authors of the study state that the etiology of [sleep problems](#) and disorders need to be identified in children with ADHD, in order to create a modified treatment regime for [sleep disorders](#) and [ADHD](#) symptoms.

Source: American Academy of Sleep Medicine ([news](#) : [web](#))

Citation: Study links ADHD with sleep problems in adolescents (2009, May 1) retrieved 15 May 2024 from <https://medicalxpress.com/news/2009-05-links-adhd-problems-adolescents.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.