

Regulating the sugar factory in diabetes

May 21 2009

Scientists in Sydney and Boston believe they may have identified a gene that controls abnormal production of sugar in the liver, a very troublesome problem for people with diabetes.

The [liver](#) is the sugar factory for the body - when blood sugar (glucose) levels fall, the liver makes and releases more. In people with diabetes, especially Type 2 diabetes, the liver doesn't stop making sugar when it should, so blood sugar levels can rise overnight while they sleep even though they haven't eaten.

Dr Jenny Gunton, diabetes specialist and endocrinologist from Sydney's Garvan Institute of Medical Research, in collaboration with Dr Xiao Hui Wang and Professor Ronald Kahn from Harvard Medical School and Joslin Diabetes Centre in Boston, have published their findings in the journal *Cell Metabolism*, now online.

"A lot of my patients will complain that they go to bed with a blood sugar of 5 and wake up with a blood sugar of 12," said Dr Gunton.

"It upsets people when their blood sugar behaves as if they're getting up in the night and having a really big snack. I have to tell them it's just one of those unfair things about having diabetes."

People with Type 2 diabetes do not produce enough [insulin](#) in the pancreas after a meal. At the same time, they are less able to use that insulin to move glucose into fat and muscle cells, a condition known as 'insulin resistance'.

With her colleagues in Boston, Gunton has been studying a transcription factor, or kind of 'master regulator', called ARNT, which controls expression of other genes involved in processes like glucose breakdown and insulin production. In an earlier study, the group showed that there is 90% less ARNT in insulin-producing cells of people with Type 2 diabetes.

The current study looks at how ARNT might be affecting the liver, and its results confirmed Dr Gunton's suspicions. "We've shown that there's likely to be decreased ARNT in the liver of people with Type 2 diabetes compared to people without Type 2 diabetes," she said.

"Working with mice, we found that [glucose levels](#) were elevated and there was glucose production from a 'precursor', a source not normally metabolised."

Other results in the study show that to some extent ARNT is regulated by insulin, so that [insulin resistance](#) in itself will contribute to a decrease in ARNT. If liver cells are treated with insulin, there will be a small increase in ARNT protein. The insulin will also help move the ARNT into the nucleus of the cell, where it does its job as a master regulator.

The paper concludes that a decline in ARNT isn't limited to the beta cells of people with [Type 2 diabetes](#). ARNT is also reduced in other important diabetes-related tissues like the liver.

Dr Gunton believes that if a new drug could be developed to increase ARNT activity in the liver, then it may be possible to shut down abnormal sugar production and improve [blood sugar](#) control in people with [diabetes](#).

Source: Research Australia ([news](#) : [web](#))

Citation: Regulating the sugar factory in diabetes (2009, May 21) retrieved 8 May 2024 from <https://medicalxpress.com/news/2009-05-sugar-factory-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.