

Implanted defibrillators: New recommendations for drivers with ICDs

June 21 2009

Patients with an Implantable Cardioverter Defibrillator (ICD) have an ongoing risk of sudden incapacitation that might cause harm to others while driving a car. Driving restrictions are imposed making these recommendations an important guideline for patients.

A consensus statement with recommendations for drivers with ICD's was presented at a press conference at the Europace 2009 meeting, in Berlin, Germany on Sunday 21 June.

A team of twelve experts from the European Heart Rhythm Association (EHRA), the Council on Cardiovascular Nursing and Allied Professions in ESC (CCNAP) and the Section [Cardiac Rehabilitation](#) of the European Association of Cardiovascular Prevention and Rehabilitation, reviewed the literature, assessed the risk and issued a consensus statement.

"[Driving](#) restrictions vary across different countries in Europe. We hope the document may serve as an instrument for European and National regulatory authorities to formulate uniform driving regulations", explained Johan Vijgen, chairperson of the task force*.

"Driving restrictions are perceived as difficult for patients and their family and have an immediate consequence for their lifestyle. In addition to the psychological and societal impact, the driving ban may also pose a considerable impact on employment and education and thereby economic status", said Vijgen.

The document presents recommendations for private driving (group 1) and professional driving (group 2). Definitions of the European Council Directives (80/1263/EEC) and (91/439/EEC) are used.

- Group 1: drivers of ordinary motor cycles, cars, and other small vehicles with or without a trailer.
- Group 2: drivers of vehicles over 3.5 metric tonnes or passenger carrying vehicles exceeding eight seats excluding the driver.

Since the introduction of the ICD in the early 1980s, multiple trials have demonstrated the efficacy of ICDs for the prevention of sudden arrhythmic death. This resulted in a significant increase in the number of implants. In Western Europe alone, 63000 ICDs were implanted in 2006 and 85500 ICDs in 2008.

Many patients are currently implanted for primary prevention (treatment of patients at risk for life-threatening arrhythmias who have never had sustained ventricular arrhythmias). The risk for sudden incapacitation is lower in these patients. Therefore, driving restriction should be less strict for these patients, than for patients implanted for secondary prevention (those who have survived a life-threatening arrhythmia).

"Patients and their families should receive adequate discharge education and standardized information on driving recommendations. This should result in a better adherence to the recommendations. It should be emphasized that the risk is mainly a consequence of the underlying condition and not of the presence of the ICD", explained Prof Vijgen.

The consensus statement will be published in the June issue of *Europace*, the official journal of the European Heart Rhythm Association.

*Members of the task force include Johan Vijgen (chairperson) Belgium, Gianluca Botto (Italy) , John Camm (United Kingdom), Carl-Johan Højer (Sweden), Werner Jung (Germany), Jean-Yves Le Heuzey (France), Andrzej Lubinski (Poland), Tone M. Norekvål (Norway), Maurizio Santomauro (Italy), Martin Schalij (The Netherlands), Jean-Paul Schmid (Switzerland), and Panos Vardas (Greece)

Source: European Society of Cardiology ([news](#) : [web](#))

Citation: Implanted defibrillators: New recommendations for drivers with ICDs (2009, June 21) retrieved 30 April 2024 from <https://medicalxpress.com/news/2009-06-implanted-defibrillators-drivers-icds.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--