

Back to normal: Surgery improves outcomes for spine patients

June 1 2009

People with the spine disease called degenerative spondylolisthesis* -- who choose surgical treatment -- experience substantially greater relief from pain over time compared to those who do not have surgery, according to a study published in the June 2009 issue of *The Journal of Bone and Joint Surgery (JBJS)*. In the past, physicians had been uncertain whether surgery provided significantly greater relief for patients, but these results help to confirm the advantages to surgery.

"There are thousands of surgeries completed each year to address degenerative spine conditions, yet, there has never been a large-scale trial to give us evidence that the surgeries really work, as compared to non-operative approaches," said study author James Weinstein, DO, MS, Third Century Professor and Chair of the departments of orthopaedics at Dartmouth Medical School and Dartmouth-Hitchcock Medical Center.

Dr. Weinstein and his colleagues collected data from 607 men and women diagnosed with spondylolisthesis who were enrolled in the Spine Patient Outcomes Research Trial (SPORT), a multi-center study that included participants from 13 medical centers in 11 states. The study was the largest ever conducted of spondylolisthesis patients.

"Until this study, our 'evidence' was anecdotal and based on patient reports. We wanted data-based, scientific evidence that we could share with patients to help them make their decisions about taking an operative vs. non-operative approach," Weinstein said.

Prior to completion of the study, SPORT looked at the three most common back conditions leading to [surgery](#), which are:

- herniated disc;
- spinal stenosis; and
- spinal stenosis with degenerative spondylolisthesis.

To be included in the study, all patients had to meet certain criteria, including:

- nerve [pain](#) in the legs
- spinal stenosis revealed on cross-sectional imaging
- degenerative spondylolisthesis evident in radiograph imaging
- symptoms which lasted for at least 12 weeks
- physician confirmation that the patient was a surgical candidate.

"Our results indicate that in these patients, there was a clear advantage for surgery," said Dr. Weinstein. "[Patients](#) felt relief faster and at two and four years, reported better function, less pain, and higher satisfaction than those who chose to go the non-surgical route."

Approximately 80 percent of Americans suffer from back pain at some point in their lives. Back pain is the most common cause of work-related disability, as well as the most expensive in terms of workers compensation and medical costs. Degenerative spondylolisthesis is one

example of this kind of painful back condition.

"Degenerative spine disease can be a debilitating condition. When well informed, surgery is a good treatment choice," said Weinstein.

Source: American Academy of Orthopaedic Surgeons ([news](#) : [web](#))

Citation: Back to normal: Surgery improves outcomes for spine patients (2009, June 1) retrieved 2 May 2024 from

<https://medicalxpress.com/news/2009-06-surgery-outcomes-spine-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--