

Give children iron supplements: They don't increase malaria risk

July 8 2009

Iron supplements do not increase the likelihood of contracting malaria and should not be withheld from children at risk of the disease, despite World Health Organisation (WHO) guidelines to the contrary, a new review by Cochrane Researchers suggests.

"Based on our review, [children](#) should not be denied iron supplements, even if they are living in areas where malaria is prevalent," says lead researcher, Juliana Ojukwu of the Department of Paediatrics at the Ebonyi State University in Ebonyi State, Nigeria. "Iron is important for growth and development, and maintaining a healthy [immune system](#)."

Until 2007, WHO guidelines recommended that all children should be given iron supplements to help prevent iron deficiency and anaemia, which are significant public health problems in developing countries. It is estimated that [iron deficiency](#) is the cause of 726,000 childhood deaths each year. However, a recent large trial in Zanzibar prompted the WHO to change its guidelines, which now recommend that iron supplements are withheld from children under two years in areas where they are at high risk of contracting malaria. The argument against giving iron is that it could help promote the growth of malarial parasites circulating in the blood.

In response to this, Cochrane researchers reviewed data from 68 different trials involving 42,981 children. They concluded that iron did not increase the risk of malaria, as long as regular [malaria](#) surveillance and treatment services were available, and that there should not be any

need to screen for anaemia before giving supplements. They say WHO guidelines rely too heavily a single recent trial, whereas this current research drew its conclusions after giving appropriate weight to a wide range of studies.

Although the benefits of giving iron are greater for children with anaemia, any decision to withhold iron supplements should be carefully considered. "Any potential negative effects of giving iron have to be weighed against the quite serious implications of not giving it, namely anaemia and its contribution to childhood infection and death, especially in Sub-Saharan Africa," says Ojukwu.

Source: Wiley ([news](#) : [web](#))

Citation: Give children iron supplements: They don't increase malaria risk (2009, July 8)
retrieved 25 April 2024 from
<https://medicalxpress.com/news/2009-07-children-iron-supplements-dont-malaria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.