

# Condoms associated with moderate protection against herpes simplex virus 2

July 13 2009

---

Condom use is associated with a reduced risk of contracting herpes simplex virus 2, according to a report based on pooled analysis of data from previous studies in the July 13 issue of *Archives of Internal Medicine*.

Herpes simplex virus 2 ([HSV-2](#)) typically causes genital [herpes](#), a chronic, life-long, viral infection. Although studies indicate that consistent condom use reduces the spread of HIV and other sexually transmitted diseases such as [chlamydia](#) and gonorrhea, the effectiveness of preventing the transmission of HSV-2 through condom use is less certain, according to background information in the article.

Emily T. Martin, M.P.H., Ph.D., of Children's Hospital Research Institute and the University of Washington, Seattle, and colleagues analyzed data from six HSV-2 studies to assess the effectiveness of condom use in preventing the virus. The studies included three candidate HSV-2 vaccine studies, an HSV-2 drug study, an observational sexually transmitted infection (STI) incidence study and a behavioral STI intervention study. These yielded results from 5,384 HSV-2-negative individuals (average age 29) at baseline for a combined total of 2,040,894 follow-up days.

More than 66 percent of those who took part in the six studies were male, 60.4 percent were white, 94.1 percent were heterosexual and most reported no prior STIs.

A total of 415 of the individuals acquired HSV-2 during follow-up. "Consistent condom users [used 100 percent of the time] had a 30 percent lower risk of HSV-2 acquisition compared with those who never used condoms," the authors write. "Risk of HSV-2 acquisition decreased by 7 percent for every additional 25 percent of the time that condoms were used during anal or vaginal sex." The risk of acquiring the virus increased significantly with increasing frequency of [unprotected sex](#) acts. There were no significant differences found in condom effectiveness between men and women.

"Based on findings of this large analysis using all available prospective data, [condom](#) use should continue to be recommended to both men and women for reducing the risk of HSV-2 acquisition," the authors conclude. "Although the magnitude of the protective effect was not as large as has been observed with other STIs, a 30 percent reduction in HSV-2 incidence can have a substantial benefit for individuals as well as a public health impact at the population level."

More information: Arch Intern Med. 2009;169[13]:1233-1240.

Source: JAMA and Archives Journals ([news](#) : [web](#))

Citation: Condoms associated with moderate protection against herpes simplex virus 2 (2009, July 13) retrieved 28 April 2024 from <https://medicalxpress.com/news/2009-07-condoms-moderate-herpes-simplex-virus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---