

High calcium level in arteries may signal serious heart attack risk

July 28 2009

Researchers may be able to predict future severe cardiac events in patients with known, stable coronary artery disease (CAD) using coronary calcium scoring, according to a study published in the online edition of *Radiology*.

"The amount of calcium in the coronary vessels, as measured by CT, is of high predictive value for subsequent serious or fatal heart attack in these patients, independent of the patient's age, sex and other coronary risk factors," said the study's lead author, Marcus Hacker, M.D., resident physician in the Department of Nuclear Medicine, leader of the research unit for nuclear cardiology and assistant medical director at Ludwig Maximilians University in Munich, Germany.

CAD is the most common type of heart disease. According to the National Heart, Lung and Blood Institute, it is the leading cause of death in the U.S. for both men and women, killing more than 500,000 Americans each year.

CAD is a condition in which plaque, consisting of cholesterol, calcium, fat and other substances, builds up inside the arteries that supply blood to the heart. When plaque builds up in the coronary arteries, blood flow to the heart is reduced and may lead to arrhythmia, heart attack or heart failure.

Single photon emission computed tomography (SPECT) myocardial perfusion imaging is a nuclear medicine diagnostic procedure that

provides excellent three-dimensional images of the coronary arteries to assist in the diagnosis and treatment of CAD.

Currently, calcium scoring—measuring the amount of calcium in the arteries—is used as a screening exam and in cases of suspected CAD, but not in cases of known CAD.

Dr. Hacker and colleagues set out to determine if calcium scoring would lend additional prognostic value to SPECT findings in patients with known, stable CAD.

For the study, 260 patients with CAD underwent coronary artery calcium scoring in addition to SPECT myocardial perfusion imaging. Over a median period of 5.4 years, the patients were followed up for severe cardiac events, meaning cardiac death or nonfatal heart attacks. Twenty-three of the 260 patients had a fatal or severe heart attack, and 40 additional patients underwent bypass surgery.

The results showed that patients with an initial calcium score greater than 400 were at significantly increased risk for severe cardiac events.

"We found that coronary calcium seems to play an important role in predicting subsequent [heart attack](#) or sudden cardiac death, and adds prognostic value to SPECT findings," said co-author Christopher Uebleis, M.D., member of the research unit for nuclear cardiology at Ludwig Maximilians University.

Dr. Hacker pointed out that combining calcium scoring and SPECT can help to identify patients with known CAD who are at highest risk for serious or fatal heart attacks.

"In these patients, intensified medical therapy, shorter follow-up intervals and, if necessary, bypass procedures may be required to prevent

future severe cardiac events."

More information: radiology.rsna.org/

Source: Radiological Society of North America ([news](#) : [web](#))

Citation: High calcium level in arteries may signal serious heart attack risk (2009, July 28)
retrieved 8 May 2024 from
<https://medicalxpress.com/news/2009-07-high-calcium-arteries-heart.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|