

High-fat, high-sugar foods alter brain receptors

July 28 2009

Overconsumption of fatty, sugary foods leads to changes in brain receptors, according to new animal research at Johns Hopkins University School of Medicine. The new research results are being presented at the 2009 annual meeting of the Society for the Study of Ingestive Behavior (SSIB), the foremost society for research into all aspects of eating and drinking behavior. The results have implications for understanding bulimia and other binge eating disorders.

Dr. Bello and colleagues report that either continuous eating or binge eating a high fat, high sugar diet alters opioid receptor levels in an area of the [brain](#) that controls [food intake](#). Opioids are a family of chemicals with actions similar to those of morphine; however, opioids exist naturally in the brain and have been linked to feelings of pleasure and euphoria. "These results are interesting because we saw changes in opioid receptor gene expression in a brain area that controls how much we eat during a meal", said Bello.

The new findings suggest that overconsumption of highly palatable foods maintains bingeing by enhancing opioids in the brain, and that increased opioids could be a factor involved in binge eating disorders. These findings may help to understand the biological basis of eating disorders.

Source: Society for the Study of Ingestive Behavior

Citation: High-fat, high-sugar foods alter brain receptors (2009, July 28) retrieved 18 April 2024 from <https://medicalxpress.com/news/2009-07-high-fat-high-sugar-foods-brain-receptors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.