

New lab test helps predict kidney damage

July 24 2009

Acute kidney injury (AKI) is a frequent complication in patients in intensive care. A new laboratory test called urine neutrophil gelatinase associated lipocalin (NGAL) helps predict if patients will develop acute kidney injury, reports an upcoming study in the *Journal of the American Society of Nephrology* (JASN). "As a stand-alone marker, urine NGAL performed moderately well in predicting ongoing and subsequent AKI," comments T. Alp Ikizler, MD (Vanderbilt University).

Another study, also in JASN, indicates that <u>urine</u> NGAL may also help in diagnosing HIV-related kidney disease affecting African Americans and black Africans. "NGAL was very specifically expressed in renal cysts—generating the new idea that NGAL may control the development of cysts in HIV-associated nephropathy," says Jonathan Barasch, MD, PhD (Columbia University, New York). He adds, "We and Prasad Devarajan, MD, identified NGAL in the kidney 10 years ago and its translation into a clinical entity in such a short time is quite a story. Almost every paper is positive for the association of NGAL and renal dysfunction/disease."

In the ICU study, patients with higher urine NGAL levels were more likely to develop acute kidney injury, even after adjustment for other factors. The rise in NGAL was present before any change in the standard test for AKI (serum creatinine level). Without other information, however, urine NGAL was no more effective in predicting AKI than clinical risk factors. Ikizler notes the study was limited by a lack of information on incidence of death or the need for dialysis, and by a lack of information on the patients' initial <u>kidney function</u> level.



In the HIV study, levels of urine NGAL were much higher in patients with HIV-associated nephropathy (HIVAN) than in patients with other forms of kidney disease, with or without HIV. HIVAN is an important complication of <u>HIV</u>, occurring mainly in patients of African descent.

Studies in mice suggested that NGAL may play an important role in the development of tubular disease and microcysts, which are specific features of HIVAN. Barasch notes that the human component of their study was limited by its small size but highlights the need for larger studies that definitively measure the NGAL monomer. He adds, "If our results are confirmed, measuring urine NGAL might help triage patients into different risk categories."

More information: The studies will appear online at jasn.asnjournals.org/ on July 23, 2009. Dr. Barasch's study is entitled, "Urinary NGAL Marks Cystic Disease in HIV-Associated Nephropathy," doi 10.1681/ASN.2009010065 . Dr. Ikizler's study is entitled, "Urine Neutrophil Gelatinase-Associated Lipocadoi 10.1681/ASN.2008070673 ute Kidney Injury in Critically III Adults," doi 10.1681/ASN.2008070673 .

Source: American Society of Nephrology (<u>news</u> : <u>web</u>)

Citation: New lab test helps predict kidney damage (2009, July 24) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2009-07-lab-kidney.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.