

Poor sleep is independently associated with depression in postpartum women

July 1 2009

A study in the July 1 issue of the journal *Sleep* suggests that postpartum depression may aggravate an already impaired sleep quality, as experiencing difficulties with sleep is a symptom of depression. Twenty-one percent of depressed postpartum women included in the study reported having also been depressed during pregnancy and 46 percent reported at least one previous depressive episode prior to conception, suggesting that new mothers diagnosed with postpartum depression are not merely reporting symptoms of chronic sleep deprivation.

Results indicate that two months after delivery, poor [sleep](#) was associated with depression when adjusted for other significant risk factors, such as poor partner relationship, previous depression, depression during [pregnancy](#) and stressful life events. [Sleep disturbances](#) and subjective sleep quality were the aspects of sleep most strongly associated with depression. Overall, nearly 60 percent of the postpartum women experienced poor global sleep quality, and 16.5 percent had depressive symptoms.

According to lead author Karen Dørheim, MD, PhD, psychiatrist at Stavanger University Hospital in Norway, depression after delivery is often not identified by new mothers, whereas tiredness and lack of sleep are common complaints. These symptoms may be attributed to poor sleep, but the tiredness could also be caused by depression.

"It is important to ask a new mother suffering from tiredness about how poor sleep affects her daytime functioning and whether there are other

factors in her life that may contribute to her lack of energy," said Dørhei. "There are also helpful depression screening questionnaires that can be completed during a consultation. Doctors and other health workers should provide an opportunity for postpartum women to discuss difficult feelings."

Data were collected between October 2005 and September 2006 from 2,830 women who gave birth to a live child at Stavanger University Hospital in Norway. Sleep was measured using the Pittsburgh Sleep Quality Index (PSQI) and depressive symptoms using the Edinburgh Postnatal Depression Scale (EPDS).

The mean self-reported nightly sleep duration was 6.5 hours, and sleep efficiency was 73 percent. The mean age of the mothers at the time of reply was 30 years, and the mean age of the infants was 8.4 weeks.

Depression, previous sleep problems, being a first time mother, not exclusively breastfeeding or having a younger or male infant were factors associated with poor postpartum [sleep quality](#). Better maternal sleep was associated with the baby sleeping in a different room.

According to authors, the first three months after delivery are characterized by continually changing sleep parameters. Women who are tired during this period may attribute this to poor sleep, but the tiredness could alternatively be caused by depression; thus talking about sleep problems may provide an entry point for also discussing the woman's overall well-being. Individual women may react differently to shorter sleep duration and lower sleep efficiency during the postpartum period, and that the sleep of women with a history of [depression](#) may be more sensitive to the psychobiological (hormonal, immunological, psychological and social) changes associated with childbirth.

Source: American Academy of Sleep Medicine ([news](#) : [web](#))

Citation: Poor sleep is independently associated with depression in postpartum women (2009, July 1) retrieved 5 May 2024 from <https://medicalxpress.com/news/2009-07-poor-independently-depression-postpartum-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.