

# Blood thinner causes stroke in some dialysis patients

August 27 2009

---

The blood thinner warfarin can prevent strokes in most individuals with abnormal heart rhythms, but the drug may have the opposite effect in kidney disease patients on dialysis, according to a study appearing in an upcoming issue of the *Journal of the American Society Nephrology* (JASN). The results suggest that warfarin should be prescribed with caution in patients with kidney failure.

People with atrial fibrillation, the most common type of [abnormal heart rhythm](#), have a significantly increased risk of [stroke](#). Blood thinners can mitigate this risk. However, almost no studies have examined whether [warfarin](#) can also prevent stroke in atrial fibrillation patients who are also on dialysis.

To investigate, Kevin Chan, MD, Michael Lazarus, MD, Raymond Hakim, MD, PhD (Fresenius Medical Care North America); and Ravi Thadhani, MD (Massachusetts General Hospital), studied 1,671 end stage renal disease (ESRD) patients with a diagnosis of atrial fibrillation who started dialysis in clinics operated by Fresenius Medical Care North America. Patients' health was monitored for an average of 1.6 years after dialysis was started.

The researchers found that patients who took warfarin increased their risk of stroke; however, the increase in risk was related to the degree of blood thinning in response to the drug. Dialysis patients whose blood was not monitored for warfarin levels after they received the drug had the highest risk of stroke (almost three-fold higher than patients not on

warfarin). Notably, clopidogrel and aspirin—two other types of blood thinners—did not increase dialysis patients' risk of stroke.

Additional research is needed to determine why warfarin has the opposite effect on stroke risk in kidney disease patients on dialysis compared with other individuals. One possible explanation may be because bleeding is a well known complication of [kidney failure](#) and the concomitant use of blood thinners may exaggerate the risk of stroke from bleeding in the head. Also, this study's findings should be validated through randomized controlled clinical trials. "Until then, physicians should be cognizant of the possible risks associated with warfarin use for atrial fibrillation in ESRD patients with careful evaluation of the risks and benefits of intervention at the individual patient level," they concluded.

More information: The article, entitled "Warfarin Use Associates with Increased Risk for Stroke in Hemodialysis Patients with [Atrial Fibrillation](#)," will appear online on August 27, 2009, [doi 10.1681/ASN.2009030319](#)

Source: American Society of Nephrology ([news](#) : [web](#))

Citation: Blood thinner causes stroke in some dialysis patients (2009, August 27) retrieved 18 April 2024 from <https://medicalxpress.com/news/2009-08-blood-thinner-dialysis-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--