

## Gallbladder emptying in primary sclerosing cholangitis patients

## August 10 2009

Primary sclerosing cholangitis (PSC) is an idiopathic chronic cholestatic inflammatory liver disease characterized by diffuse fibrosing inflammation of intra- and/or extrahepatic bile ducts, resulting in bile duct obliteration, biliary cirrhosis, and eventually hepatic failure. One of the most common symptoms at the time of presentation of PSC is mild to severe abdominal pain localized in the right upper quadrant. However, the mechanisms responsible for the abdominal pain in PSC are not fully understood.

A research team led by Karouk Said from Karolinska University Hospital addressed this question. Their study will be published on July 28, 2009 in the *World Journal of Gastroenterology*.

In this study, 20 patients with PSC and 10 healthy subjects were investigated. Gallbladder fasting volume, ejection fraction and residual volume after ingestion of a test meal were compared in patients with PSC and healthy controls using magnetic resonance imaging. Symptoms, thickness and contrast enhancement of the gallbladder wall and the presence of cystic duct strictures were also assessed.

Median fasting gallbladder volume in patients with PSC [67 (19-348) mL] was twice that in healthy controls [32 (16-55) mL] (P

This concluded that PSC patients have increased fasting gallbladder volume. Gallbladder mucosal dysfunction secondary to chronic cholecystitis, may be a possible mechanism for increased gallbladder



volume. Gallbladder size or emptying does not seem to be involved in the development of <u>abdominal pain</u> in patients with PSC.

More information: Said K, Edsborg N, Albiin N, Bergquist A. Gallbladder emptying in patients with primary sclerosing cholangitis. *World J Gastroenterol* 2009; 15(28): 3498-3503 www.wjgnet.com/1007-9327/15/3498.asp

Source: World Journal of Gastroenterology (news: web)

Citation: Gallbladder emptying in primary sclerosing cholangitis patients (2009, August 10) retrieved 25 April 2024 from

https://medicalxpress.com/news/2009-08-gallbladder-primary-sclerosing-cholangitis-patients.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.