


'Achilles' heel' in Y chromosome linked to sex disorders

September 3 2009



Isodicentric Y chromosomes are generated during sperm production through aberrant crossing over between opposite arms of a palindrome. At left is two copies of the normal Y chromosome, each carrying the masculinizing gene SRY and a centromere. Also shown is one of the Y chromosome's palindromes, depicted here as a pair of blue and red arrowheads pointing in opposite directions. The big "X" denotes a "crossing over" between opposing arms of the same palindrome on the two Y chromosomes. The result is the isodicentric Y chromosome diagrammed at right. The isodicentric has two copies of SRY, two centromeres, and is a mirror-image structure. Credit: Tom DiCesare/Whitehead Institute

The unique mechanism behind the evolutionary survival of the human Y chromosome may also be responsible for a range of sex disorders, from failed sperm production to sex reversal to Turner Syndrome.




Roughly six years ago, David Page's lab at Whitehead Institute for Biomedical Research reported the discovery of eight large areas of mirror-imaged genetic sequences, or palindromes, along the Y chromosome. Because the Y chromosome essentially has no partner with which to swap genes, a process that between ordinary chromosome pairs leads to genetic diversity and the exchange of good genes for damaged ones, it relies on its own palindromes to swap genes with itself. The Y, as it turns out, folds itself in the middle of palindromic regions, thereby pairing identical sequences to allow for potentially beneficial genetic exchange.

At the time, the finding provided explanation for why, despite much-heralded reports to the contrary, the Y chromosome is not doomed to extinction. Now, the Page lab has discovered that the Y's process of self-preservation can randomly go awry, with considerable clinical consequence.

"This is the sequel to the Y chromosome palindrome story," says Page, Whitehead Institute Director and a Howard Hughes Medical Institute investigator.

The latest chapter of the story, whose conclusion is published in the September 4 issue of *Cell*, began with the intriguing hypothesis that perhaps the Y's process of self-recombination can inadvertently turn the entire chromosome into a palindrome—literally, a mirror-image of itself. The result would be a so-called isodicentric Y chromosome (idicY), an abnormal structure with, as the name implies, two centromeres.





When a fertilized human egg carries an isodicentric Y chromosome, the likelihood of sex reversal (anatomic feminization) during fetal development depends on the size of the isodicentric. Diagrammed here are four different isodicentric Y chromosomes, each with two copies of the masculinizing gene SRY and two centromeres. Among these four isodicentrics, the distance between the two centromeres, and hence the size of the isodicentric, increases from left to right. The intensity of the mitotic tug of war (previous figure), and hence the likelihood of loss of the dicentric, increases in the same direction. Mitotic loss of the isodicentric in fetal gonads causes sex reversal (anatomic feminization), whose likelihood increases with the size of the isodicentric. Credit: Image: Tom DiCesare/Whitehead Institute


"We began to think seriously about the centromeres and the activity around them. Two centromeres render the chromosome susceptible to damage," says Julian Lange, first author of the *Cell* paper and a former graduate student in the Page lab. Because of the Y chromosome's well known roles in sex determination and male fertility, Lange began to speculate about the potential clinical impact of the transmission of an idicY during fertilization.

"Because the Y chromosome is not essential to an individual's survival, these isodicentric Ys can persist," says Lange, who, after completing this research at Whitehead, became a postdoctoral fellow at Memorial Sloan-



Kettering Cancer Center. "They can be found in the population."

And Lange found them, in the DNA samples of 51 patients screened from a field of nearly 2400 individuals who had come under study over the course of many years because of failed sperm production, structurally abnormal Y chromosomes, or sex reversal. Through sophisticated genetic analysis, it became clear that idicYs were responsible for spermatogenic failure in many of the male patients.



A tug of war between nascent daughter cells can snap isodicentric in two, leading to its loss during development of the fetus. Credit: Tom DiCesare/Whitehead

However, 18 of the 51 patients were anatomically female—despite having two copies of the male-determining SRY gene on their idicY chromosomes. Lange and Page began to hypothesize that the feminization they had identified was related to the instability of the idicYs themselves. Suspecting that the level of instability of an idicY increases with the size of the chromosome, they looked for a connection and found something somewhat paradoxical: the larger the Y chromosome, the greater the likelihood of sex reversal.



"We had predicted this correlation, which relates to the overall distance between the centromeres," says Page. "But when we confirmed it with the patient data, we were blown away."

Page believes that this new model for the formation of idicY chromosomes, coupled with the size-instability correlation, suggests a causal link to Turner syndrome, a chromosomal abnormality in girls or women, characterized by the lack of one sex chromosome. Turner syndrome affects an estimated 1 in 2500 females. Page won't yet speculate as to what percentage of Turner syndrome could be caused by this palindrome-to-palindrome recombination, but he does think it's significant.

Source: Whitehead Institute for Biomedical Research

Citation: 'Achilles' heel' in Y chromosome linked to sex disorders (2009, September 3) retrieved 10 April 2024 from

https://medicalxpress.com/news/2009-09-achilles-heel-chromosome-linked-sex.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.